

## Eleventh exercise sheet on Relativity and Cosmology I

Winter term 2018/19

**Release:** Mon, Jan. 14<sup>th</sup>

**Submit:** Mon, Jan. 21<sup>st</sup> in lecture

**Discuss:** Thu, Jan. 24<sup>th</sup>

### Exercise 35 (10 points): *An exact plane-wave solution I*

Consider the following *exact* metric given by Bondi, Pirani and Robinson (1959):

$$ds^2 = L^2 \left( e^{+2\beta} dx^2 + e^{-2\beta} dy^2 \right) + dz^2 - dt^2 = L^2 \left( e^{+2\beta} dx^2 + e^{-2\beta} dy^2 \right) - du dv,$$

where  $L = L(u)$ ,  $\beta = \beta(u)$  are the *background* and *wave factors*, and  $u = t - z$ ,  $v = t + z$  the *retarded* and *advanced light-cone* or *null coordinates*, respectively.

**35.1** Show that

$$\begin{aligned} \Gamma^v_{xx} &= \frac{d}{du} \left( L^2 e^{+2\beta} \right) & \Gamma^v_{yy} &= \frac{d}{du} \left( L^2 e^{-2\beta} \right); \\ \Gamma^x_{ux} = \Gamma^x_{xu} &= \frac{1}{2} \frac{d}{du} \ln \left( L^2 e^{+2\beta} \right) & \Gamma^y_{uy} = \Gamma^y_{yu} &= \frac{1}{2} \frac{d}{du} \ln \left( L^2 e^{-2\beta} \right), \end{aligned}$$

whereas all the other Christoffel symbols of the second kind vanishes.

*Hint.* It might be quicker to apply the variational method directly, instead of using the definition.

**35.2** Show that a *co-moving* particle, defined by  $\dot{x} = \dot{y} = \dot{z} = 0$  and  $\dot{t} = 1$ , moves on a geodesic.

**35.3** Derive the Ricci tensor.

### Exercise 36 (6+7 points): *An exact plane-wave solution II*

The vacuum Einstein equation of the Bondi–Pirani–Robinson metric reduces to

$$L'' + (\beta')^2 L = 0, \quad f' := \frac{df}{du}.$$

**36.1** Solve the vacuum Einstein equation for  $\beta' \equiv 0$ ,  $L(0) = 1$  and  $L'(0) = 0$ .

**36.2** For simplicity, let  $\beta \equiv 0$ ,  $L(0) = 1$  and  $L'(0) = -1$ . Solve the vacuum Einstein equation for these conditions. Employ the transformation to the new coordinates

$$\begin{aligned} \bar{t} &= t - \frac{1}{2}(1 - t + z) \left( x^2 + y^2 \right), \\ \bar{z} &= z - \frac{1}{2}(1 - t + z) \left( x^2 + y^2 \right), \\ \bar{x} &= (1 - t + z)x, \quad \bar{y} = (1 - t + z)y \end{aligned}$$

to show that this solution is in fact just the Minkowski space, although containing a singularity.

**36.3** Consider the linearised theory. For that purpose let  $\beta$  and  $\beta'$  be small and of the same order.

Show that the vacuum Einstein equations can be integrated, such that the result corresponds to a  $+$ -polarised wave in the linearised theory.

**36.4** (bonus) Now go back to the full theory. Let  $L \equiv 1$  for  $u \leq -a < 0$ , and  $\beta' \equiv 0$  for  $u < -a$  and  $u \geq 0$ . In other words,  $\beta' \neq 0$  can only happen for  $-a \leq u < 0$ . Furthermore, assume that  $\beta'$  is such that the *singularity* appears at some  $u > 0$ .

Provide a *qualitative* discussion and a sketch of the solution for  $L(u)$ .

*Hint:* It is sufficient to consider the convexity of  $L$  on  $[-a, 0)$ .

**36.5** (bonus) Recall the configuration in the lecture, where a beam of plane gravitational wave meets a ring of test particles, initially at rest in the  $z = 0$  plane.

What happens to the ring after the wave has passed by, in the exact **(36.4)** and the linearised theories?

### Exercise 37 (4+3 points): *Polarisation*

In a flat background spacetime, consider two Cartesian coordinate systems  $(t, x, y, z)$  and  $(t, x', y', z)$  that can be transformed into each other by a rotation with the angle  $\theta$  around the  $z$ -axis.

**37.1** Consider an electromagnetic wave that propagates in the  $z$ -direction. Let  $\hat{\mathbf{e}}_x, \hat{\mathbf{e}}_y, \hat{\mathbf{e}}_{x'},$  and  $\hat{\mathbf{e}}_{y'}$  be the unit polarisation vectors in the coordinate systems. Show that

$$\hat{\mathbf{e}}_{x'} = \hat{\mathbf{e}}_x \cos(\theta) + \hat{\mathbf{e}}_y \sin(\theta), \quad \hat{\mathbf{e}}_{y'} = -\hat{\mathbf{e}}_x \sin(\theta) + \hat{\mathbf{e}}_y \cos(\theta).$$

**37.2** Analogously, consider a linearised gravitational wave propagating in the  $z$ -direction. Let  $\mathbf{e}_+, \mathbf{e}_x, \mathbf{e}_{+'},$   $\mathbf{e}_{x'}$  be the polarisation tensors in the coordinate systems. Show that

$$\mathbf{e}_{+'} = \mathbf{e}_+ \cos(2\theta) + \mathbf{e}_x \sin(2\theta), \quad \mathbf{e}_{x'} = -\mathbf{e}_+ \sin(2\theta) + \mathbf{e}_x \cos(2\theta).$$

**37.3** (bonus) Let  $|\rightarrow\rangle$  and  $|\leftarrow\rangle$  be the quantum states of a spin- $\frac{1}{2}$  particle, whose spin is aligned or anti-aligned with respect to the  $x$ -direction, respectively, and analogously  $|\rightarrow'\rangle$  and  $|\leftarrow'\rangle$  with respect to the  $x'$ -direction. Show that

$$|\rightarrow'\rangle = |\rightarrow\rangle \cos\left(\frac{\theta}{2}\right) + i|\leftarrow\rangle \sin\left(\frac{\theta}{2}\right), \quad |\leftarrow'\rangle = i|\rightarrow\rangle \sin\left(\frac{\theta}{2}\right) + |\leftarrow\rangle \cos\left(\frac{\theta}{2}\right).$$

**37.4** (bonus) Write down the generalisation for the basis states of linear polarisation for a radiation field of arbitrary spin  $s$ .