

## 1<sup>st</sup> exercise sheet on Relativity and Cosmology II

Summer term 2019

**Release:** Mon, Apr. 1<sup>st</sup>

**Submit:** Mon, Apr. 8<sup>th</sup> in lecture

**Discuss:** Thu, Apr. 11<sup>th</sup>

### Exercise 41 (20 = 10 + 6 + 4 points): *Effective Schwarzschild potential*

The aim of this exercise is to analyse certain properties of the movement of massive test particles in the Schwarzschild space-time.

For this purpose, consider the equation of motion on the equatorial plane  $\theta = \pi/2$  with an effective potential  $V_{\text{eff}}$  that results from the geodesic equation

$$\frac{\dot{r}^2}{2} + V_{\text{eff}}(r) = E, \quad V_{\text{eff}}(r) = -\frac{GM}{r} + \frac{\ell^2}{2r^2} - \frac{GM\ell^2}{r^3}.$$

Here  $\ell$  and  $E$  indicate constants of motion.

In the following, express radial distances in terms of the Schwarzschild radius  $r_S = 2GM$ .

- 41.1**
- Analyse and sketch the potential  $V_{\text{eff}}(r)$  for all relevant cases (characterised by the values of  $M$  and  $\ell$ ).
  - In which cases do bound particle orbits exist? Analyse the stability of all orbits.
  - Which conditions does a test particle approaching from infinity ( $r \rightarrow +\infty$ ) have to fulfil in order to fall into the centre of the effective potential?  
Under which circumstances does a particle that starts from rest at infinity fall into the centre?
  - Compare the results obtained so far to the situation in Newtonian gravity.
  - Show the following statements:
    - For  $\ell/GM < 2\sqrt{3}$  every in-falling particle falls towards the event horizon  $r = 2GM$ .
    - The most strongly bound orbit is located at  $r = 6GM$  with  $\ell/GM = 2\sqrt{3}$  and it possesses a relative binding energy of  $1 - \sqrt{8/9}$ .

**41.2** Consider a massive test particle initially being at rest at the radial coordinate  $R > r_S$  that falls radially ( $\ell = 0$ ) into the centre.

- Find the solution of the resulting initial value problem.  
*Hint:* The solution can be given in a parametrised form  $r(\eta)$ ,  $\tau(\eta)$  (where  $\tau$  is the proper time of the particle) which describes a *cycloid* orbit.  
At which proper time  $\tau_0$  does the particle reach the centre of the potential?
- How long does it take for this particle to reach the Schwarzschild radius as measured by an observer at infinity?

**41.3** In the lecture it was mentioned that Kepler's Third Law

$$GM = \omega^2 r^3, \quad \text{where } \omega = d\phi/dt,$$

holds for circular orbits in the Schwarzschild space-time. Prove this statement.