

Second exercise sheet on Relativity and Cosmology II

Summer term 2021

Release: Mon, Apr. 26th

Submit: Mon, May 3th on ILIAS

Discuss: Thr, May 6th

In the following exercises, consider the coordinates (t, r, θ, ϕ) . The 2-dimensional equatorial spatial slice Σ is defined by $t = \text{const.}$, $\theta = \pi/2$, and its induced metric reads $d\sigma_\Sigma^2$. Moreover, the Schwarzschild metric in such coordinates reads

$$ds^2 = -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega^2, \quad f(r) := 1 - \frac{R_S}{r}, \quad R_S := 2GM, \quad d\Omega^2 := d\theta^2 + \sin^2 \theta d\phi^2.$$

Furthermore, denote the Euclidean metric on \mathbb{R}^n by $d\vec{x}_{\mathbb{R}^n}^2$.

Exercise 38 (10 points): *Schwarzschild metric in isotropic coordinates*

Consider a coordinate transformation

$$r = \left(1 + \frac{R_S}{4\bar{r}}\right)^2 \bar{r}, \quad t = \bar{t}, \quad \theta = \bar{\theta}, \quad \phi = \bar{\phi}.$$

38.1 Express the Schwarzschild metric in the new coordinates $(\bar{t}, \bar{r}, \bar{\theta}, \bar{\phi})$, which are called *isotropic*.

38.2 Compare briefly the metric components in old and new coordinates at the event horizon $R_S = 2GM$.

38.3 In the *standard* coordinates, consider a radial range $R_S < r < R$.

Use the *isotropic* coordinates to calculate

- the surface area on the equatorial spatial slice between these radii, and
- the volume of a spherical shell with $t = \text{const.}$ within the range.

38.4 Compare your results in **38.3** to the corresponding quantities in the Euclidean space.

Exercise 39 (4 points): *Isometric embedding I: the Schwarzschild space*

39.1 Consider the cylindrical coordinates (ρ, ψ, z) of \mathbb{R}^3 . Set

$$d\vec{x}_{\mathbb{R}^3}^2 \Big|_{z=z(r)} \equiv d\sigma_\Sigma^2$$

and integrate the resulting equation (*Flamm's paraboloid*).

39.2 In the current case, if $d\sigma_\Sigma^2 \rightarrow d\vec{x}_{\mathbb{R}^2}^2$ as $r \rightarrow +\infty$, Σ is called *asymptotically flat*.

Analytically extend the embedding in **39.1**, and show that there can be *two* distinct regions, which are both asymptotically flat (*Einstein and Rosen*).

Hint: Sketching the embedding may help.

Exercise 40 (6 points): *Isometric embedding II: a wormhole*

Consider a spacetime \mathcal{M}_W with the metric

$$ds^2 = -dt^2 + dr^2 + (b^2 + r^2)d\Omega^2$$

where b is a constant of dimension length.

40.1 Argue briefly that the equatorial spatial slice Σ of \mathcal{M}_W is representative for the latter.

40.2 Consider the cylindrical coordinates (ρ, ψ, z) of \mathbb{R}^3 . Set

$$d\vec{x}_{\mathbb{R}^3}^2 \Big|_{\rho=\rho(r), z=z(r)} \equiv d\sigma_{\Sigma}^2$$

and integrate the resulting equation for z and ρ .

40.3 Argue briefly that there is a hole-like structure in \mathcal{M}_W .

Hint: Sketching the embedding may help.

Remark. The Einstein tensor of the metric reads

$$G_{\mu\nu} dx^\mu dx^\nu = \frac{b^2}{(b^2 + r^2)^2} (-dt^2 - dr^2 + (b^2 + r^2)d\Omega^2).$$

If the matter were modelled by an ideal fluid, its energy density would turn out to be negative,

$$\rho = T_{\mu\nu} u^\mu u^\nu = -\frac{1}{\kappa} \frac{b^2}{(b^2 + r^2)^2} < 0.$$

In other words, matter with negative energy density is needed to source this spacetime.

Inversely, one may also ask, if it is possible to have a viable wormhole spacetime sourced by matter with positive energy density. Under precise and additional assumptions, this has been excluded by the so-called *topological censorship theorem*.