

8th exercise sheet on Relativity and Cosmology I

Winter term 2015/16

Deadline for delivery: Thursday, 17th December 2015 during the exercise class.

Exercise 21 (6 credit points): *Integration*

Let us define a surface of revolution \mathcal{A} via

$$z^2 = [f(r)]^2, \quad r^2 = x^2 + y^2,$$

where f is a strictly positive smooth function on $[0, a)$ with $f(a) = f'(0) = 0$ and $f'(a) = -\infty$.

21.1 Determine the metric g_{ij} which is induced by the line element $ds^2 = dr^2 + r^2 d\varphi^2 + dz^2$ on this surface.

21.2 Calculate the corresponding Ricci scalar and show by explicit calculation that the integral

$$\int_{\mathcal{A}} \sqrt{g} R d^2x$$

does not depend on the choice of the function f .

Exercise 22 (14 credit points): *Killing vector fields*

22.1 Show that the Killing equation $\nabla_{(\mu} v_{\nu)} = 0$ can also be written as $\mathcal{L}_v g_{\mu\nu} = 0$.

What does this mean from a physical point of view?

22.2 Give a proof for the following integrability condition for a Killing vector field v^μ :

$$v_{\lambda;\kappa\nu} = -v_\mu R^\mu{}_{\nu\lambda\kappa}.$$

22.3 Consider a timelike Killing vector field ξ^μ . Show that there is a coordinate system in which the metric does not depend on time, i.e. $\frac{\partial g_{\mu\nu}}{\partial t} = 0$ holds.

22.4 Find all Killing vector fields for Minkowski spacetime.

22.5 Let $u^\mu = dx^\mu/d\tau$ be the tangent vector of a geodesic, i.e. $u^\nu \nabla_\nu u^\mu = 0$, and let ξ^μ be a Killing vector field. Show that $u_\mu \xi^\mu$ is constant along the geodesic. Use these conserved quantities to illustrate the physical meaning of the Killing vector fields from part **23.4**.

22.6 Let $T^{\mu\nu}$ be a symmetric tensor field with vanishing divergence, i.e. $\nabla_\mu T^{\mu\nu} = 0$, and let ξ^μ be a Killing vector field. Calculate $(\xi^\mu T_\mu{}^\nu)_{;\nu}$.

The result is of great importance for the construction of conserved integral quantities.