

8th exercise sheet on Relativity and Cosmology II

Summer term 2014

Deadline for delivery: Wednesday, 4th June 2014 during the exercise class.

Exercise 15 (16 credit points): *Kerr–Newman metric*

The most general solution for a stationary black hole is given by the *Kerr–Newman metric*, which describes a black hole with angular momentum $J = Ma$ and charge q . The line element expressed in Boyer–Lindquist coordinates takes the following form:

$$ds^2 = -\frac{\Delta}{\rho^2} \left(dt - a \sin^2(\theta) d\phi \right)^2 + \frac{\sin^2(\theta)}{\rho^2} \left[(r^2 + a^2) d\phi - a dt \right]^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2,$$

where

$$\rho^2 = r^2 + a^2 \cos^2(\theta), \quad \Delta = r^2 - 2Mr + q^2 + a^2, \quad q^2 + a^2 \leq M^2.$$

15.1 Show that this line element arises from the line element of the Kerr metric by means of the substitution $M \rightarrow M - q^2/(2r)$.

15.2 For $\Delta = 0$ the metric exhibits coordinate singularities. Determine their radial coordinates r_{\pm} .

The surface $r_+ = \text{const.}$ (with r_+ being the radial coordinate with a larger value) represents the event horizon. Calculate its surface area for $t = \text{const.}$

15.3 Analogously to the Kerr metric, consider an observer with $r = \text{const.}$, $\theta = \pi/2$, whose tangent vector is parallel to the Killing field $\chi^\mu = \zeta^\mu + \Omega \Psi^\mu$.

Which values can Ω take for given $r \geq r_+$? Show that at the horizon only one value Ω_H is possible and determine this value.

15.4 Consider the Killing field $\chi^\mu = \zeta^\mu + \Omega \Psi^\mu$ evaluated at the event horizon.

Show that this Killing field is light-like on the entire horizon. Furthermore, show that the surface gravity κ defined by means of $[\nabla^\mu(\chi_\nu \chi^\nu)]_H = -2\kappa \chi^\mu|_H$ is a well-defined quantity.

Calculate the Lie derivative of the defining equation for κ with respect to χ^μ and thereby show that κ is constant along the integral curves of χ .

Remark: After a rather long calculation one obtains $\kappa = (r_+ - M)/(r_+^2 + a^2)$. (Not to be shown here.)

15.5 Consider the null geodesics defined at the horizon, whose tangent vectors k^μ are proportional to χ^μ .

Find the functional relationship between the affine parameter λ of these null geodesics and the Killing parameter v of the integral curves of χ^μ , i.e. $\chi^\mu = (\partial/\partial v)^\mu$.

Exercise 16 (4 credit points): *Hawking temperature*

In the lecture it was mentioned that a Schwarzschild black hole radiates with the so-called *Hawking temperature*

$$T_H = \frac{\hbar c^3}{8\pi k_B G M}.$$

Assume that only photons are emitted and that they have a perfect Planck spectrum. Find a relation between the initial mass of the black hole and its lifetime and analyze this relation for several interesting masses and time intervals.