

School on quantum many-body
phenomena out of equilibrium
August 25/26, 2023
ICTP Trieste, Italy



Driven Open Quantum Systems: From Micro- to Macrophysics

Sebastian Diehl

Institute for Theoretical Physics, University of Cologne

Outline

Keldysh theory general: A. Kamenev, *Field theory of non-equilibrium systems*, Cambridge University Press

Review: L. Sieberer, M. Buchhold, SD, *Keldysh Field Theory for Driven Open Quantum Systems*, Reports on Progress in Physics (2016)


Lecture I & II: Theoretical background & non-equilibrium phases

- From the quantum master equation to the Keldysh functional integral
 - construction
 - semiclassical limit, connection to exciton-polariton systems
 - “what is non-equilibrium about it?”
- Applications: stationary states of driven open quantum systems
 - fate of BKT physics out of equilibrium
 - phase transition induced by non-equilibrium drive

$$\partial_t \rho = -i[H, \rho] + \mathcal{L}[\rho]$$




$$e^{i\Gamma[\Phi]} = \int \mathcal{D}\delta\Phi e^{iS_M[\Phi+\delta\Phi]}$$




Lecture III: Lindblad-Keldysh 2.0: Measurement induced phase transitions of fermions

- weak continuous measurements
- replica approach and replica Keldysh functional integral
- BKT transition of replica fluctuations




Lindblad quantum master equation: From few to many degrees of freedom



$$\partial_t \rho = -i[H, \rho] + \kappa \sum_i L_i \rho L_i^\dagger - \frac{1}{2} \{L_i^\dagger L_i, \rho\}$$


What is a driven open quantum system?

- quantum optics:



- exchange** between system and bath (e.g. energy, entropy, particle number)

- example: laser driven atom coupled to the radiation field (two-level system)



- simple fact: **drive essential** to access upper level

- implications:

- no** minimisation of **energy**
- no** guarantee for detailed balance
- no** obedience of the **second law** of thermodynamics (state purification)

Driven open quantum systems: microscopic description


- quantum master equation

$$\partial_t \hat{\rho} = \underbrace{-i[\hat{H}, \hat{\rho}]}_{\text{coherent evolution}} + \sum_i \underbrace{\gamma_i [2\hat{L}_i \hat{\rho} \hat{L}_i^\dagger - \hat{L}_i^\dagger \hat{L}_i \hat{\rho} - \hat{\rho} \hat{L}_i^\dagger \hat{L}_i]}_{\text{driven-dissipative evolution}}$$

Lindblad operators


lattice site, spin ...

$\equiv \hat{\mathcal{L}}[\hat{\rho}]$ Lindbladian; also: Liouvillian



- derivation from system-bath setting: second order time dependent perturbation theory

- example: two-level system



$$H = (|e\rangle, |g\rangle) \begin{pmatrix} \Delta & \Omega \\ \Omega & 0 \end{pmatrix} \begin{pmatrix} \langle e| \\ \langle g| \end{pmatrix} \quad \hat{L} = |g\rangle\langle e| = \sigma^-$$

$g_\mu \sim \Omega$

- starting point: system-bath setting

$$\hat{H}_b = \sum \epsilon_\mu \hat{b}_\mu^\dagger \hat{b}_\mu$$

$$\hat{H}_t = \hat{H} + \hat{H}_b + \hat{H}_{s-b}$$

$$\hat{H}_{s-b} = \sum_\mu g_\mu e^{-i\nu t} \hat{L} \hat{b}_\mu^\dagger + \text{h. c.}$$

- bath: infinitely many mode modes than system (for single L)

- 3 approximations:

- Born**: weak system-bath coupling -> bath unaffected by system (2nd order pert. th.)

- Markov**: system evolution slow wrt bath -> time-local evolution


- rotating wave**: drive ν selects relevant energy regimes

Driven open quantum systems: microscopic description

- quantum master equation

$$\partial_t \hat{\rho} = \underbrace{-i[\hat{H}, \hat{\rho}]}_{\text{coherent evolution}} + \sum_i \underbrace{\gamma_i [2\hat{L}_i \hat{\rho} \hat{L}_i^\dagger - \hat{L}_i^\dagger \hat{L}_i \hat{\rho} - \hat{\rho} \hat{L}_i^\dagger \hat{L}_i]}_{\text{driven-dissipative evolution}}$$

Lindblad operators



The diagram shows a light blue rounded rectangle labeled 'environment' containing a red oval labeled 'system'. A green arrow labeled 'drive' points into the system from the right.

$$\equiv \hat{\mathcal{L}}[\hat{\rho}] \quad \text{Lindbladian; also: Liouvillian}$$

- derivation from ‘symmetry’ (i.e. implementing key physical requirements)

- Lindbladian defines a **dynamical map** $\hat{\rho}(t + \Delta t) = \hat{\rho}(t) + \Delta t \cdot \hat{\mathcal{L}}[\hat{\rho}]$
- with properties

- Hermiticity: $\hat{\rho}(t)^\dagger = \hat{\rho}(t) \implies \hat{\rho}^\dagger(t + \Delta t) = \hat{\rho}(t + \Delta t)$ since $\hat{\mathcal{L}}[\hat{\rho}]^\dagger = \hat{\mathcal{L}}[\hat{\rho}]$

- complete positivity: $\hat{\rho}(t) \geq 0 \implies \hat{\rho}(t + \Delta t) \geq 0$

- trace preservation / probability conservation $\partial_t \text{tr} \hat{\rho}(t) = 0$ since $\text{tr} \hat{\mathcal{L}}[\hat{\rho}] = 0$

→ up to a unitary transformation (above: diagonal form in index i), $\hat{\mathcal{L}}[\hat{\rho}]$ is the most general **time-local generator** with these properties

G. Lindblad, *Commun. Math. Phys.* (1976)

Nielsen & Chuang, Chap. 8




Göran Lindblad

Driven open quantum systems: microscopic description

- quantum master equation

$$\partial_t \hat{\rho} = \underbrace{-i[\hat{H}, \hat{\rho}]}_{\text{coherent evolution}} + \sum_i \underbrace{\gamma_i [2\hat{L}_i \hat{\rho} \hat{L}_i^\dagger - \hat{L}_i^\dagger \hat{L}_i \hat{\rho} - \hat{\rho} \hat{L}_i^\dagger \hat{L}_i]}_{\text{driven-dissipative evolution}}$$

Lindblad operators



The diagram shows a light blue rounded rectangle labeled 'environment' containing a red oval labeled 'system'. A green arrow labeled 'drive' points from the right into the 'system' oval.

$$\equiv \hat{\mathcal{L}}[\hat{\rho}] \quad \text{Lindbladian; also: Liouvillian}$$

- interpretation:

$$\partial_t \hat{\rho} = \underbrace{-i\hat{H}}_{\text{energy}} - \underbrace{i \sum_i \gamma_i \hat{L}_i^\dagger \hat{L}_i}_{\text{decay (dissipation)}} \hat{\rho} + \text{h.c.} + \underbrace{2 \sum_i \gamma_i \hat{L}_i \hat{\rho} \hat{L}_i^\dagger}_{\substack{\text{ensures probability conservation} \\ \text{(fluctuation)}}}$$

“ $E - i\Gamma$ ”


$\partial_t \text{tr} \hat{\rho}(t) = 0$

Driven open quantum systems: microscopic description

- quantum master equation

$$\partial_t \hat{\rho} = \underbrace{-i[\hat{H}, \hat{\rho}]}_{\text{coherent evolution}} + \sum_i \underbrace{\gamma_i [2\hat{L}_i \hat{\rho} \hat{L}_i^\dagger - \hat{L}_i^\dagger \hat{L}_i \hat{\rho} - \hat{\rho} \hat{L}_i^\dagger \hat{L}_i]}_{\text{driven-dissipative evolution}}$$

Lindblad operators



The diagram shows a light blue rounded rectangle labeled 'environment' containing a red oval labeled 'system'. A green arrow labeled 'drive' points into the system from the right.

$$\equiv \hat{\mathcal{L}}[\hat{\rho}] \quad \text{Lindbladian; also: Liouvillian}$$

- So far: **few degrees of freedom** in the “system”
- Question: What if we replace few by **many degrees of freedom**?

➡ The interface of quantum optics and many-body physics

➡ **Quantum Optics:**
coherent and driven-dissipative
dynamics on equal footing

➡ **Many-Body Physics:**
continuum of spatial
degrees of freedom

➡ **Statistical Mechanics:**
physics at the largest
distances

microphysics

macrophysics

The interface of quantum optics and many-body physics

➔ Quantum Optics

➔ Many-Body Physics


➔ Statistical Mechanics

microphysics

macrophysics

- The experimental platforms: light-matter systems realize **driven open quantum matter**


Atoms



Bose-Einstein condensate in a cavity

Baumann et al., Nature (2010)


Light



Microcavity arrays

Houck, Türeci, Koch, Nat. Phys. (2012)

Solids




Exciton-polariton condensates

Kasprzak et al., Nature (2006)

and more:


- polar molecules
- nano-mechanics
- photon BECs



driven-dissipative Rydberg gases


S. Helmrich, A. Arias, G. Lochhead, M. Buchhold, SD, S. Whitlock, Nature (2020); T. Wintermantel, ... SD, S. Whitlock, Nat. Comm. (2021)

Quantum devices / NISQ Platforms




superconducting circuits

K. Satzinger et al. Science (2021)



Rydberg tweezers

G. Semeghini et al. Science (2021)



trapped ions

C. Noel et al. Nat. Phys. (2022)

The interface of quantum optics and many-body physics

➡ Quantum Optics

➡ Many-Body Physics

➡ Statistical Mechanics

microphysics

Microscopic


“Thermodynamic”

macrophysics

Long wavelength

- Questions and challenges to theory: physics at various length scales

Novel universal phenomena ?




Efficient theoretical tools ?

$$Z[J] = \int \mathcal{D}\varphi e^{i(S[\varphi] + \int J\varphi)}$$

perform the transition from micro-to
macrophysics:

quantum field theory out of equilibrium

Experimental platforms ?



cold atoms, light-driven semiconductors, microcavity
arrays, trapped ions, NISQ ...

A workhorse model: Lindbladian formulation

- generic microscopic many-body model: 'Lindblad ϕ^4 theory'

$$\partial_t \rho = -i[H, \rho] + \mathcal{D}[\rho] \equiv \mathcal{L}[\rho]$$

$$H = \int_{\mathbf{x}} \hat{\phi}_{\mathbf{x}}^\dagger \left(\frac{\Delta}{2M} - \mu \right) \hat{\phi}_{\mathbf{x}} + \frac{\lambda}{2} (\hat{\phi}_{\mathbf{x}}^\dagger \hat{\phi}_{\mathbf{x}})^2$$

$$\mathcal{D}[\rho] = \underbrace{\gamma_p \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}}^\dagger \rho \hat{\phi}_{\mathbf{x}} - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}} \hat{\phi}_{\mathbf{x}}^\dagger, \rho \}]}_{\text{single particle pump}} + \underbrace{\gamma_l \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}} \rho \hat{\phi}_{\mathbf{x}}^\dagger - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}}^\dagger \hat{\phi}_{\mathbf{x}}, \rho \}]}_{\text{single particle loss}} +$$

$$\underbrace{\kappa \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}}^2 \rho \hat{\phi}_{\mathbf{x}}^{\dagger 2} - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}}^{\dagger 2} \hat{\phi}_{\mathbf{x}}^2, \rho \}]}_{\text{two particle loss}}$$

- basic physics: mean field theory



- study evolution of $\phi(\mathbf{x}, t) \equiv \langle \hat{\phi}(\mathbf{x}) \rangle(t) = \text{tr}[\hat{\phi}(\mathbf{x}) \hat{\rho}(t)]$ $\hat{\rho} = \prod_{\mathbf{x}} \hat{\rho}(\mathbf{x}), \quad \hat{\rho}(\mathbf{x}) = |\phi(\mathbf{x})\rangle \langle \phi(\mathbf{x})|$

coherent state

- homogenous limit $\phi(\mathbf{x}, t) = \phi(t)$

$$i\partial_t \phi(t) = [-\mu - i(\gamma_l - \gamma_p) + (\lambda - i\kappa)|\phi(t)|^2]\phi(t)$$

- overdamped motion in potential landscape
- condensation / spontaneous U(1) symmetry breaking for $\gamma_l - \gamma_p < 0$




A workhorse model: Lindbladian formulation

- generic microscopic many-body model: ‘Lindblad ϕ^4 theory’

$$\partial_t \rho = -i[H, \rho] + \mathcal{D}[\rho] \equiv \mathcal{L}[\rho]$$


$$H = \int_{\mathbf{x}} \hat{\phi}_{\mathbf{x}}^\dagger \left(\frac{\Delta}{2M} - \mu \right) \hat{\phi}_{\mathbf{x}} + \frac{\lambda}{2} (\hat{\phi}_{\mathbf{x}}^\dagger \hat{\phi}_{\mathbf{x}})^2$$

$$\begin{aligned} \mathcal{D}[\rho] = & \underbrace{\gamma_p \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}}^\dagger \rho \hat{\phi}_{\mathbf{x}} - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}} \hat{\phi}_{\mathbf{x}}^\dagger, \rho \}]}_{\text{single particle pump}} + \underbrace{\gamma_l \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}} \rho \hat{\phi}_{\mathbf{x}}^\dagger - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}}^\dagger \hat{\phi}_{\mathbf{x}}, \rho \}]}_{\text{single particle loss}} + \\ & \underbrace{\kappa \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}}^2 \rho \hat{\phi}_{\mathbf{x}}^{\dagger 2} - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}}^{\dagger 2} \hat{\phi}_{\mathbf{x}}^2, \rho \}]}_{\text{two particle loss}} \end{aligned}$$



- plan:

- translate to Lindblad-Keldysh functional integral



- how does this model relate e.g. to exciton-polariton systems? (semiclassical limit)
- ‘what is non-equilibrium about it’?
- how to extract the phase structure?



Leonid W. Keldysh

Keldysh functional integral for stationary states of driven open quantum systems

- Construction from quantum master equation
- Semiclassical limit
- “What is non-equilibrium about it?”



$$\partial_t \hat{\rho} = -i(\hat{H} - \sum_i \gamma_i \hat{L}_i^\dagger \hat{L}_i) \hat{\rho} + \text{h.c.} + 2 \sum_i \gamma_i \hat{L}_i \hat{\rho} \hat{L}_i^\dagger$$

$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])}$$

Keldysh functional integrals: Why?

- Feynman's formulation of quantum mechanics



Space-Time Approach to Non-Relativistic Quantum Mechanics

R. P. FEYNMAN

Cornell University, Ithaca, New York

Non-relativistic quantum mechanics is formulated here in a different way. It is, however, mathematically equivalent to the familiar formulation. In quantum mechanics the probability of an event which can happen in several different ways is the absolute square of a sum of complex contributions, one from each alternative way. The probability that a particle will be found to have a path $x(t)$ lying somewhere within a region of space time is the square of a sum of contributions, one from each path in the region. The contribution from a single path is postulated to be an exponential whose (imaginary) phase is the classical action (in units of $\hbar$) for the path in question. The total contribution from all paths reaching x, t from the past is the wave function $\psi(x, t)$. This is shown to satisfy Schroedinger's equation. The relation to matrix and operator algebra is discussed. Applications are indicated, in particular to eliminate the coordinates of the field oscillators from the equations of quantum electrodynamics.

1. INTRODUCTION

IT is a curious historical fact that modern quantum mechanics began with two quite different mathematical formulations: the differential equation of Schroedinger, and the matrix algebra of Heisenberg. The two, apparently dissimilar approaches, were proved to be mathematically equivalent. These two points of view were destined to complement one another and to be ultimately synthesized in Dirac's transformation theory.

This paper will describe what is essentially a

classical action³ to quantum mechanics. A probability amplitude is associated with an entire motion of a particle as a function of time, rather than simply with a position of the particle at a particular time.

The formulation is mathematically equivalent to the more usual formulations. There are, therefore, no fundamentally new results. However, there is a pleasure in recognizing old things from a new point of view. Also, there are problems for which the new point of view offers a distinct advantage. For example, if two systems

- Useful language for systems with **many degrees of freedom**
- general: powerful techniques
 - diagrammatic perturbation theory;
 - collective variables;
 - renormalization group
- non-equilibrium Keldysh
- closer to the real-time formulations of quantum mechanics
- gives unified view on and principles (e.g. symmetries) for equilibrium and non-equilibrium systems
- indispensable for many systems:
 - disorder infinite harmonic
 - dissipation baths!
- open the powerful toolbox of quantum field theory for many-body non-equilibrium situations

Keldysh functional integral

more details: L. Sieberer, M. Buchhold, SD,
Keldysh Field Theory for Driven Open Quantum Systems,
Reports on Progress in Physics (2016)

- The basic idea in three steps:

$$\hbar = 1$$

$$U(t, t_0) = e^{-iH(t-t_0)}$$

1. Schrödinger equation: evolving a state **vector**

$$i\partial_t |\psi\rangle(t) = H|\psi\rangle(t) \quad \Rightarrow \quad |\psi\rangle(t) = U(t, t_0)|\psi\rangle(t_0)$$

2. Heisenberg-von Neumann equation: evolving a state (density) **matrix**

$$\partial_t \rho(t) = -i[H, \rho(t)] \quad \Rightarrow \quad \rho(t) = U(t, t_0)\rho(t_0)U^\dagger(t, t_0)$$

- identical for pure (separable) states $\rho = |\psi\rangle\langle\psi|$

3. The same is true for the Lindblad equation:


$$\partial_t \rho = -i[H, \rho] + \kappa \sum_i L_i \rho L_i^\dagger - \frac{1}{2} \{L_i^\dagger L_i, \rho\} \equiv \mathcal{L}[\rho]$$

$$\Rightarrow \rho(t) = e^{\mathcal{L}(t-t_0)} \rho(t_0)$$

linear **superoperator** (acts from both
sides on density matrix)

1. Functional integral idea

- ➔ “Trotterization” of time interval and insertion of coherent states: $e^{iH(t-t_0)} = \lim_{N \rightarrow \infty} (1 + i\delta_t H)^N$



- one time step: evaluate matrix element (observe structure, details later!)

$$e^{-\phi_n^* \phi_n} \langle \phi_{n+1} | e^{-i\delta_t H[a^\dagger, a]} | \phi_n \rangle$$

$$e^{-\phi_{n+1}^* \phi_{n+1}} | \phi_{n+1} \rangle \langle \phi_{n+1} | e^{-\phi_n^* \phi_n} | \phi_n \rangle \langle \phi_n |$$

$e^{-i\delta_t H}$

$$\approx e^{idt[-i\phi(t)\partial_t \phi(t)^* - H[\phi^*(t), \phi(t)]]}$$

$$\delta_t \rightarrow 0$$

- many time steps:

$$\underbrace{\int \prod_t \frac{d\phi^*(t) d\phi(t)}{\pi}}_{=: \int \mathcal{D}(\phi^*, \phi) \text{ functional integral measure}} e^{i \underbrace{\int_{t_0}^{t_f} dt [-i\partial_t \phi^*(t) \cdot \phi(t) - H[\phi^*(t), \phi(t)]]}_{\text{action}}}$$

- ➔ operator H -> complex, time dependent functional H

coherent states (bosons):

$$a|\phi\rangle = \phi|\phi\rangle$$

$$\langle \phi' | \phi \rangle = e^{\phi'^* \phi}$$

$$\mathbf{1} = \int \frac{d\phi^* d\phi}{\pi} e^{-\phi^* \phi} |\phi\rangle \langle \phi|$$

2. Schrödinger vs. Heisenberg-von Neumann


- Schrödinger equation: evolving a state **vector** $U(t, t_0) = e^{-iH(t-t_0)}$

$$i\partial_t |\psi\rangle(t) = H|\psi\rangle(t) \Rightarrow |\psi\rangle(t) = U(t, t_0)|\psi\rangle(t_0)$$

- Heisenberg-von Neumann equation: evolving a state (density) **matrix**

$$\partial_t \rho(t) = -i[H, \rho(t)] \Rightarrow \rho(t) = U(t, t_0)\rho(t_0)U^\dagger(t, t_0)$$

- Second case: “Trotterization” on both sides:



$$e^{iH(t-t_0)} = \lim_{N \rightarrow \infty} (1 + i\delta_t H)^N \quad \delta_t = \frac{t-t_0}{N}$$

➔ **two** sets of degrees of freedom for **matrix** evolution

3. Schrödinger vs. Lindblad

- Schrödinger equation: evolving a state **vector**


$$U(t, t_0) = e^{-iH(t-t_0)}$$

$$i\partial_t |\psi\rangle(t) = H|\psi\rangle(t) \quad \Rightarrow \quad |\psi\rangle(t) = U(t, t_0)|\psi\rangle(t_0)$$

- Lindblad equation: evolving a state (density) **matrix**

$$\partial_t \rho = -i[H, \rho] + \mathcal{D}[\rho] \equiv \mathcal{L}[\rho] \quad \Rightarrow \quad \rho(t) = e^{\mathcal{L}(t-t_0)} \rho(t_0)$$

- Identical program for Liouville generator of dynamics (left and right action on density matrix)



$$\rho(t) = e^{(t-t_0)\mathcal{L}} \rho_0 = \lim_{N \rightarrow \infty} (1 + \delta_t \mathcal{L})^N \rho_0 \quad \delta_t = \frac{t - t_0}{N}$$

➔ **two** sets of degrees of freedom for **matrix** evolution

Keldysh partition function

- Schrödinger equation: evolving a state **vector**

$$U(t, t_0) = e^{-iH(t-t_0)}$$

$$i\partial_t |\psi\rangle(t) = H|\psi\rangle(t) \quad \Rightarrow \quad |\psi\rangle(t) = U(t, t_0)|\psi\rangle(t_0)$$

- Lindblad equation**: evolving a state (density) **matrix**


$$\partial_t \rho = -i[H, \rho] + \mathcal{D}[\rho] \equiv \mathcal{L}[\rho] \quad \Rightarrow \quad \rho(t) = e^{\mathcal{L}(t-t_0)} \rho(t_0)$$

- final step: Keldysh “partition function”

$$Z = \text{tr} \rho(t) = \text{tr} \rho(t_0) = 1$$



Detailed calculation: damped harmonic oscillator




- key steps present for single degree of freedom, e.g.

$$\hat{\mathcal{L}}[\rho] = -i[\omega_0 \hat{a}^\dagger \hat{a}, \rho] + \kappa(2\hat{a}\rho\hat{a}^\dagger - \{\hat{a}^\dagger \hat{a}, \rho\}) \quad \rho(t) = e^{(t-t_0)\mathcal{L}} \rho_0 = \lim_{N \rightarrow \infty} (1 + \delta_t \mathcal{L})^N \rho_0$$

$$\delta_t = \frac{t - t_0}{N}$$

- one time step



coherent states (bosons):

$$a|\phi\rangle = \phi|\phi\rangle$$

$$\langle\phi'|\phi\rangle = e^{\phi'^*\phi}$$

$$\mathbf{1} = \int \frac{d\phi^* d\phi}{\pi} e^{-\phi^*\phi} |\phi\rangle\langle\phi|$$

- many time steps in temporal continuum limit

- from a single free to many interacting degrees of freedom

- single d.o.f. $\sigma = \pm$ $\phi_\sigma(t), \int dt$ (0+1) dimensional field theory
- many d.o.f.s, lattice $\phi_{\sigma,i}(t), \int dt \sum_i$
- many d.o.f.s, continuum $\phi_\sigma(t, \vec{x}), \int dt d^d x$ (d+1) dimensional field theory

- works analogously for interactions (subtlety: operator ordering for non-linear L)
- left out: fermions (additional sign in odd-parity Lindblad operators)

Keldysh functional integral for interacting many-body system

- Lindblad equation:

$$\begin{aligned}\partial_t \rho &= -i[H, \rho] + \mathcal{D}[\rho] \\ &= -i(H\rho - \rho H) + \kappa \sum_i (L_i \rho L_i^\dagger - \frac{1}{2} L_i^\dagger L_i \rho - \frac{1}{2} \rho L_i^\dagger L_i)\end{aligned}$$

- equivalent Keldysh functional integral:

$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])} \quad \Phi_\pm = \begin{pmatrix} \phi_\pm \\ \phi_\pm^* \end{pmatrix}$$


$$S_M[\Phi_+, \Phi_-] = \int dt (\phi_+^* i \partial_t \phi_+ - \phi_-^* i \partial_t \phi_- - i \mathcal{L}[\Phi_+, \Phi_-])$$

$$\mathcal{L}[\Phi_+, \Phi_-] = -i (H_+ - H_-) - \kappa \sum_i \left(L_{i,+} L_{i,-}^\dagger - \frac{1}{2} L_{i,+}^\dagger L_{i,+} - \frac{1}{2} L_{i,-}^\dagger L_{i,-} \right)$$

$$H_\pm = H(\Phi_\pm) \text{ etc.}$$

- ➔ recognize Lindblad structure
- ➔ simple translation table (for contour normal ordered Lindbladian)

- operator right of density matrix -> - contour
- operator left of density matrix -> + contour



- ➔ caveat: contour diagonal Lindblad terms need temporal regularisation to track operator ordering

Keldysh functional integral: structural properties

- Keldysh functional integral:

$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])} \quad \Phi_{\pm} = \begin{pmatrix} \phi_{\pm} \\ \phi_{\pm}^* \end{pmatrix}$$


$$S_M[\Phi_+, \Phi_-] = \int dt (\phi_+^* i \partial_t \phi_+ - \phi_-^* i \partial_t \phi_- - i \mathcal{L}[\Phi_+, \Phi_-])$$

$$\mathcal{L}[\Phi_+, \Phi_-] = -i (H_+ - H_-) - \kappa \sum_i \left(L_{i,+} L_{i,-}^{\dagger} - \frac{1}{2} L_{i,+}^{\dagger} L_{i,+} - \frac{1}{2} L_{i,-}^{\dagger} L_{i,-} \right)$$

$$H_{\pm} = H(\Phi_{\pm}) \text{ etc.}$$

- focus on three key aspects, reflecting different levels of inclusion of fluctuations

1. Probability conservation (zero order)
2. Deterministic limit (first order)
3. Fluctuations (quantum and class. statistical) fully included



- next lecture, compromise: semiclassical limit, classical statistical fluctuations included

classical path
(minimizes S)

1. Probability conservation / "causality"

- trace / probability conservation:

- Lindblad:
$$\partial_t \text{tr} \rho = \text{tr} \left(-i(H\rho - \rho H) + \kappa \sum_i (L_i \rho L_i^\dagger - \frac{1}{2} L_i^\dagger L_i \rho - \frac{1}{2} \rho L_i^\dagger L_i) \right) = 0$$
 cyclicity

- Keldysh: $Z = \text{tr} \rho(t) = 1 \quad Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])}$


- will argue: reflected on the action as

$$\Phi_+ = \Phi_- \Rightarrow S_M[\Phi_+, \Phi_-] = 0$$

- redundancy

$$\langle \hat{O} \rangle(t) = \text{tr}[\hat{O} \hat{\rho}(t)] = \text{tr}[\hat{\rho}(t) \hat{O}]$$

$$\langle \hat{O} \rangle(t) = \langle O_+(t) \rangle = \langle O_-(t) \rangle$$



$$\implies 0 = \partial_t Z = i \langle s(\Phi_+(t), \Phi_-(t)) \rangle = i \langle s(\Phi_+(t), \Phi_+(t)) \rangle \quad \forall t \quad S = \int_{t_0}^t dt' s(\Phi_+(t'), \Phi_-(t'))$$

- mnemonic: taking trace = ignoring contour order
- motivates **Keldysh rotation**

1. Probability conservation and Keldysh rotation

- trace / probability conservation:

- Lindblad: $\partial_t \text{tr} \rho = \text{tr} \left(-i(H\rho - \rho H) + \kappa \sum_i (L_i \rho L_i^\dagger - \frac{1}{2} L_i^\dagger L_i \rho - \frac{1}{2} \rho L_i^\dagger L_i) \right) = 0$ cyclicity
- Keldysh: $Z = \text{tr} \rho(t) = 1$ $Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])}$

- make probability conservation more handy:

- starting point: **contour basis**: $S[\Phi_+, \Phi_-]$ with $S[\Phi_+, \Phi_- = \Phi_+] = 0$

- **Keldysh rotation** $\begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_+ + \phi_- \\ \phi_+ - \phi_- \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} \phi_+ \\ \phi_- \end{pmatrix}$ center-of-mass
relative

- action in **Keldysh/RAK basis** $S[\Phi_c, \Phi_q]$

- probability conservation

$$S[\Phi_c, \Phi_q = 0] = 0$$

- interpretation of the fields: use $\langle \hat{\phi} \rangle = \langle \phi_+ \rangle = \langle \phi_- \rangle$

- “**classical**” field can acquire expectation value
($\longleftrightarrow$ condensation, spontaneous symmetry breaking)

$$\langle \hat{\phi} \rangle = \langle \phi_c \rangle / \sqrt{2}, \quad \langle \phi_q \rangle = 0$$

- “**quantum**” field cannot

2. Deterministic limit

- Keldysh functional integral

$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])}$$

- probability conservation: zero order in the quantum field $S[\Phi_c, \Phi_q = 0] = 0$

- first order in quantum field: ordering principle due to **condensation**

- classical / occupation field: **macroscopic occupation** $N \rightarrow \infty$ (more precisely: only $q=0$ mode scales)


$$\phi_c(q) \sim N^{1/2} \implies \phi_c(x) \sim \frac{N^{1/2}}{V^{1/2}} \sim N^0$$

$$\phi_c(x) = V^{-1/2} \sum_q e^{-iqx} \phi_c(q)$$

- quantum field:

$$\phi_q(q) \sim N^0 \implies \phi_q(x) \sim \frac{1}{V^{1/2}} \sim N^{-1/2}$$



- expand the action to leading order $S[\Phi_+ = (\Phi_c + \Phi_q)/\sqrt{2}, \Phi_- = (\Phi_c - \Phi_q)/\sqrt{2}] \approx \int_{\mathbf{x}, t} [\phi_q \frac{\delta S}{\delta \phi_c} + \phi_q^* \frac{\delta S}{\delta \phi_c^*}]$

- can do integral over quantum field $Z = \int D[\phi_c, \phi_c^*] \delta \left[\frac{\delta S}{\delta \phi_c} \right] \delta \left[\frac{\delta S}{\delta \phi_c^*} \right]$ 

- only deterministic configuration contributes with $\frac{\delta S}{\delta \phi_c} = 0 = \frac{\delta S}{\delta \phi_c^*}$

→ deterministic limit Lindblad ϕ^4 : dissipative Gross-Pitaevskii mean field theory

exercise: make connection to operator mean field theory





2. Deterministic limit

- Keldysh functional integral

$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])}$$

- probability conservation: zero order in the quantum field

$$S[\Phi_c, \Phi_q = 0] = 0$$



- Summary:

- deterministic limit of Keldysh functional integral dominated by single field configuration minimizing bare Keldysh action

$$\frac{\delta S}{\delta \phi_c} = 0 = \frac{\delta S}{\delta \phi_c^*}$$

- applications: systems in the presence of condensation/ collective degrees of freedom and low noise level, e.g.
 - Bose condensation: dynamics of zero temperature weakly interacting Bose gases (cold atoms)
 - Light condensation: classical optics, waveguides, ... (“non-Hermitian physics”)

3. Fluctuations


- **Classical** field theories: single configuration has it all, the one minimising $S[\phi]$ i.e. $\frac{\delta S}{\delta \phi} = 0$ **deterministic**

- **Quantum /statistical** field theories: Summation over all possible field configurations

$$\int \mathcal{D}\phi e^{iS[\phi]}$$

- Keldysh functional integral

$$\int \mathcal{D}(\Phi_+, \Phi_-) e^{iS[\Phi_+, \Phi_-]}$$



- probability conservation: zero order in the quantum field $S[\Phi_c, \Phi_q = 0] = 0$
- deterministic limit: first order in quantum field
- how to quantify deviations from the deterministic limit?

- ➔ **correlation functions**: fluctuations around deterministic configuration $\sim \langle \phi_c^*(\vec{x}, t) \phi_c(\vec{x}', t') \rangle$
- det.: $= \langle \phi_c^*(\vec{x}, t) \rangle \langle \phi_c(\vec{x}', t') \rangle$
- ➔ **response functions**: impact of an external perturbation $\sim \langle \phi_q^*(\vec{x}, t) \phi_c(\vec{x}', t') \rangle$

exercise: verify relation to operator formalism!

3. Fluctuations: Correlation vs. response functions

more details: L. Sieberer, M. Buchhold, SD, Reports on Progress in Physics (2016)

- introduce **complex contour dependent** sources (cf. StatMech)

$$Z[j] = \langle e^{i \int (j_+ \phi_+^* - j_- \phi_-^* + c.c.)} \rangle = \langle e^{i \int (j_c \phi_q^* + j_q \phi_c^* + c.c.)} \rangle \quad Z[j=0] = Z = 1$$

normalization

- order parameter / occupation field:

$$\langle \phi_c(t, \mathbf{x}) \rangle = -i \frac{\delta Z[j]}{\delta j_q^*(t, \mathbf{x})} \Big|_{j=0}$$

- single particle response: how does the field react to external perturbations?

relation to operator formalism
(once and for all)

$$G^R(t-t', \mathbf{x}-\mathbf{x}') = i \frac{\delta^2 Z}{\delta j_q^*(t, \mathbf{x}) \delta j_c(t', \mathbf{x}')} \Big|_{j=0} = -i \langle \phi_c(t, \mathbf{x}) \phi_q^*(t', \mathbf{x}') \rangle \stackrel{\downarrow}{=} -i \theta(t-t') \langle [\hat{\phi}(t, \mathbf{x}), \hat{\phi}^\dagger(t', \mathbf{x}')] \rangle \stackrel{\downarrow}{=} 1$$

$t=t', \mathbf{x}=\mathbf{x}'$

- single particle correlations: how strong are fluctuations?

$$G^K(t-t', \mathbf{x}-\mathbf{x}') = i \frac{\delta^2 Z}{\delta j_q^*(t, \mathbf{x}) \delta j_q(t', \mathbf{x}')} \Big|_{j=0} = -i \langle \phi_c(t, \mathbf{x}) \phi_c^*(t', \mathbf{x}') \rangle = -i \langle \{\hat{\phi}(t, \mathbf{x}), \hat{\phi}^\dagger(t', \mathbf{x}')\} \rangle \stackrel{\downarrow}{=} 2 \langle \hat{n}(\mathbf{x}) \rangle + 1$$

$t=t', \mathbf{x}=\mathbf{x}'$

time and space translation
invariance assumed

extra: how are states occupied?

- total Green's function

$$G = \begin{pmatrix} G^K & G^R \\ G^A & 0 \end{pmatrix}$$

prob. conservation

$$G^A = (G^R)^\dagger, \quad (G^K)^\dagger = -G^K$$

Hermitian conjugates anti-Hermitian

3. Fluctuations: Correlation vs. response functions & Keldysh action

- by example: master equation for decaying cavity

more details: L. Sieberer, M. Buchhold, SD,
Reports on Progress in Physics (2016)

$$\partial_t \rho = -i[\omega_0 \hat{a}^\dagger \hat{a}, \rho] + \kappa(2\hat{a}\rho\hat{a}^\dagger - \{\hat{a}^\dagger \hat{a}, \rho\})$$

- action:

$$\begin{aligned}
 S &= \int dt (\phi_c^*, \phi_q^*) \begin{pmatrix} 0 & i\partial_t - \omega_0 - i\kappa \\ i\partial_t - \omega_0 + i\kappa & 2i\kappa \end{pmatrix} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} && \text{time domain} \\
 &&& \phi_\nu(t) \\
 &&& \equiv P^A(\omega) \\
 &= \int dt (\phi_c^*, \phi_q^*) \underbrace{\begin{pmatrix} 0 & \omega - \omega_0 - i\kappa \\ \omega - \omega_0 + i\kappa & 2i\kappa \end{pmatrix}}_{\substack{\equiv P^R(\omega) \quad \equiv P^K \\ \text{claim: } G^{-1}(\omega)}} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} && \text{frequency domain} \\
 &&& \phi_\nu(\omega)
 \end{aligned}$$

3. Fluctuations: Correlation vs. response functions & Keldysh action

- by example: master equation for decaying cavity

more details: L. Sieberer, M. Buchhold, SD,
Reports on Progress in Physics (2016)


$$\partial_t \rho = -i[\omega_0 \hat{a}^\dagger \hat{a}, \rho] + \kappa(2\hat{a}\rho\hat{a}^\dagger - \{\hat{a}^\dagger \hat{a}, \rho\})$$

- action:

$$S = \int dt (\phi_c^*, \phi_q^*) \begin{pmatrix} 0 & i\partial_t - \omega_0 - i\kappa \\ i\partial_t - \omega_0 + i\kappa & 2i\kappa \end{pmatrix} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} \quad \begin{array}{l} \text{time domain} \\ \phi_\nu(t) \end{array}$$

$$= \int dt (\phi_c^*, \phi_q^*) \underbrace{\begin{pmatrix} 0 & \omega - \omega_0 - i\kappa \\ \omega - \omega_0 + i\kappa & 2i\kappa \end{pmatrix}}_{\equiv P^R(\omega)} \underbrace{\begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix}}_{\equiv P^K} \quad \begin{array}{l} \text{frequency domain} \\ \phi_\nu(\omega) \end{array}$$

$G^{-1}(\omega)$



- partition function: Gaussian integration

$$Z[j_c, j_q] = \langle e^{i \int \frac{d\omega}{2\pi} (j_c^* \phi_q + j_q^* \phi_c + \text{h.c.})} \rangle = e^{i \int \frac{d\omega}{2\pi} (j_q^*, j_c^*) G(\omega) \begin{pmatrix} j_q \\ j_c \end{pmatrix}}$$

- single particle Green's function

$$\begin{pmatrix} \langle \phi_c(\omega) \phi_c^*(\omega') \rangle & \langle \phi_c(\omega) \phi_q^*(\omega') \rangle \\ \langle \phi_q(\omega) \phi_c^*(\omega') \rangle & \langle \phi_q(\omega) \phi_q^*(\omega') \rangle \end{pmatrix} = - \left(\begin{array}{cc} \frac{\delta^2 Z}{\delta j_q^*(\omega) \delta j_q(\omega')} & \frac{\delta^2 Z}{\delta j_q^*(\omega) \delta j_c(\omega')} \\ \frac{\delta^2 Z}{\delta j_c^*(\omega) \delta j_q(\omega')} & \frac{\delta^2 Z}{\delta j_c^*(\omega) \delta j_c(\omega')} \end{array} \right) \Big|_{j=0} = iG(\omega) \delta(\omega - \omega')$$

- summary in matrix components (generally valid — beyond Gaussian single mode example):

$$G^{-1} = \begin{pmatrix} 0 & P^A \\ P^R & P^K \end{pmatrix} \implies G = \begin{pmatrix} G^K & G^R \\ G^A & 0 \end{pmatrix} \quad G^{R/A} = [P^{R/A}]^{-1} \quad G^K = -G^R P^K G^A$$

action matrix kernel

single particle Green's function

3. Fluctuations: Correlation vs. response functions & Keldysh action

- by example: master equation for decaying cavity

more details: L. Sieberer, M. Buchhold, SD,
Reports on Progress in Physics (2016)

$$\partial_t \rho = -i[\omega_0 \hat{a}^\dagger \hat{a}, \rho] + \kappa(2\hat{a}\rho\hat{a}^\dagger - \{\hat{a}^\dagger \hat{a}, \rho\})$$

- action:

$$S = \int dt (\phi_c^*, \phi_q^*) \begin{pmatrix} 0 & i\partial_t - \omega_0 - i\kappa \\ i\partial_t - \omega_0 + i\kappa & 2i\kappa \end{pmatrix} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} \quad \begin{array}{l} \text{time domain} \\ \phi_\nu(t) \end{array}$$

$$= \int dt (\phi_c^*, \phi_q^*) \begin{pmatrix} 0 & \omega - \omega_0 - i\kappa \\ \omega - \omega_0 + i\kappa & 2i\kappa \end{pmatrix} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} \quad \begin{array}{l} \text{frequency domain} \\ \phi_\nu(\omega) \end{array}$$

- observables from the Green's functions:

-  decay of **single-particle response**: $G^R(t - t') = \int_\omega e^{i\omega(t-t')} G^R(\omega) = \theta(t - t') e^{i\omega_0(t-t')} e^{-\kappa(t-t')}$
- Lorentzian spectral density: $A(\omega) = \text{Im} G^R(\omega) = \frac{2\kappa}{(\omega - \omega_0)^2 + \kappa^2}$
-  cavity mode **occupation** in stationary state $2\langle \hat{n}(t) \rangle + 1 = \langle \hat{a}^\dagger(t) \hat{a}(t) + \hat{a}(t) \hat{a}^\dagger(t) \rangle = iG^K(t - t) = i \int_\omega e^{i\omega(t-t)} G^K(\omega) = 1$
 $\langle \hat{n}(t \rightarrow \infty) \rangle = 0 \quad (t \rightarrow \infty)$

→ correlation / statistical properties:

G^K

→ response / spectral properties:

G^R

Driven Open Quantum Systems: From Micro- to Macrophysics Lecture II

- From the quantum master equation to the Keldysh functional integral
 - construction
 - semiclassical limit, connection to exciton-polariton systems
 - “what is non-equilibrium about it?”
- Applications: stationary states of driven open quantum systems
 - fate of BKT physics out of equilibrium
 - phase transition driven by non-equilibrium drive

$$\partial_t \rho = -i[H, \rho] + \mathcal{L}[\rho]$$



$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{iS[\Phi_+, \Phi_-]}$$

microphysics



macrophysics

Back to many-body model: Workhorse Lindbladian


- generic microscopic many-body model:

$$\partial_t \rho = -i[H, \rho] + \mathcal{D}[\rho] \equiv \mathcal{L}[\rho]$$

$$H = \int_{\mathbf{x}} \underbrace{\hat{\phi}_{\mathbf{x}}^\dagger \left(\frac{\Delta}{2M} - \mu \right) \hat{\phi}_{\mathbf{x}}}_{\text{kinetic energy}} + \underbrace{\frac{\lambda}{2} (\hat{\phi}_{\mathbf{x}}^\dagger \hat{\phi}_{\mathbf{x}})^2}_{\text{two-particle interaction}}$$

$$\mathcal{D}[\rho] = \underbrace{\gamma_p \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}}^\dagger \rho \hat{\phi}_{\mathbf{x}} - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}} \hat{\phi}_{\mathbf{x}}^\dagger, \rho \}]}_{\text{single particle pump}} + \underbrace{\gamma_l \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}} \rho \hat{\phi}_{\mathbf{x}}^\dagger - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}}^\dagger \hat{\phi}_{\mathbf{x}}, \rho \}]}_{\text{single particle loss}} +$$

$$\underbrace{\kappa \int_{\mathbf{x}} [\hat{\phi}_{\mathbf{x}}^2 \rho \hat{\phi}_{\mathbf{x}}^{\dagger 2} - \frac{1}{2} \{ \hat{\phi}_{\mathbf{x}}^{\dagger 2} \hat{\phi}_{\mathbf{x}}^2, \rho \}]}_{\text{two-particle loss}}$$



Many-Body Master Equation

1-1
mapping

Keldysh functional
integral






Many-body model: Workhorse Lindblad-Keldysh action

- after Keldysh rotation:

$$\mathcal{S} = \int_{t,\mathbf{x}} \left\{ (\phi_c^*, \phi_q^*) \underbrace{\begin{pmatrix} 0 & P^A \\ P^R & P^K \end{pmatrix}}_{\text{after Fourier: } G^{-1}(\omega, \mathbf{q})} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} + 2i\kappa \phi_c^* \phi_c \phi_q^* \phi_q - \frac{1}{2} [(\lambda + i\kappa) (\phi_c^{*2} \phi_c \phi_q + \phi_q^{*2} \phi_c \phi_q) + c.c.] \right\}$$

after Fourier: $G^{-1}(\omega, \mathbf{q})$



- Gaussian sector: inverse Green's function

- retarded/advanced $P^R(\omega, \mathbf{q}) = \omega - \frac{(\mathbf{q}^2 - \mu)}{2M} + i(\gamma_l - \gamma_p)/2$

- Keldysh component $P^K = i(\gamma_l + \gamma_p)$

difference: distance from a phase transition


sum: noise of loss and pumping add up

- now: simplifications in the semiclassical limit:

- sharp argument close to a critical point
- provides intuition for a frequency regime $\omega \ll \gamma = \gamma_l + \gamma_p$


- now: “what is non-equilibrium about it?”

Semi-classical limit and Langevin equations




Intermezzo: Exciton-polariton systems

Kasprzak et al., Nature 2006



Imamoglu et al., PRA 1996



- phenomenological description: stochastic driven-dissipative Gross-Pitaevskii-Eq

$$i\partial_t\phi = \left[\underbrace{-\frac{\nabla^2}{2m}}_{\text{propagation}} - \mu + \underbrace{i(\gamma_p - \gamma_l)}_{\text{pump \& loss rates}} + \underbrace{(\lambda - i\kappa)}_{\substack{\text{elastic collisions} \\ \text{two-body loss}}} |\phi|^2 \right] \phi + \zeta$$

Szymanska, Keeling, Littlewood PRL (04,06); PRB (07);
Wouters, Carusotto PRL (07,10)

$$\langle \zeta^*(t, \mathbf{x}) \zeta(t', \mathbf{x}') \rangle = \gamma \delta(t - t') \delta(\mathbf{x} - \mathbf{x}')$$


- structure:
$$i\partial_t\phi = \underbrace{\frac{\delta\mathcal{H}_c}{\delta\phi^*}}_{\text{coherent (reversible)}} + \underbrace{i\frac{\delta\mathcal{H}_d}{\delta\phi^*}}_{\text{dissipative (irreversible)}} + \underbrace{\zeta}_{\substack{\text{noise} \leftrightarrow \\ \text{random force allowing to explore} \\ \text{various configurations}}}$$

$$\mathcal{H}_\alpha = \int d^3x [r_\alpha |\phi|^2 + K_\alpha |\nabla\phi|^2 + \lambda_\alpha |\phi|^4]$$

$$\alpha = c, d$$

Intermezzo: Exciton-polariton systems

- Bose condensation seen despite non-equilibrium conditions



Kasprzak et al., Nature 2006


- stochastic driven-dissipative Gross-Pitaevskii-Eq

~~$$i\partial_t \phi = \left[-\frac{\nabla^2}{2m} - \mu + i(\gamma_p - \gamma_l) + (\lambda - i\kappa) |\phi|^2 \right] \phi + \xi$$~~

Szymanska, Keeling, Littlewood PRL (04, 06); PRB (07)); Wouters, Carusotto PRL (07,10)

- mean field

- neglect noise
- homogeneous solution $\phi(\mathbf{x}, t) = \phi_0$




- naively, just as Bose condensation in equilibrium!
- Q1: How does this model relate to the Lindbladian and Lindblad-Keldysh field theory?
- Q2: What is “non-equilibrium” about it?

Semiclassical limit of Lindblad-Keldysh action: power counting

$$\mathcal{S} = \int_{t,\mathbf{x}} \left\{ (\phi_c^*, \phi_q^*) \begin{pmatrix} 0 & P^A \\ P^R & P^K \end{pmatrix} \begin{pmatrix} \phi_c \\ \phi_q \end{pmatrix} + 2i\kappa \phi_c^* \phi_c \phi_q^* \phi_q - \frac{1}{2} [(\lambda + i\kappa) (\phi_c^{*2} \phi_c \phi_q + \phi_q^{*2} \phi_c \phi_q) + c.c.] \right\}$$

- Gaussian sector **close to a critical point**:



- retarded/advanced $P^R(\omega, \mathbf{q}) = \omega - \mathbf{q}^2 - \mu + i(\gamma_l - \gamma_p)/2 \sim q^2$

- Keldysh component $P^K = i(\gamma_l + \gamma_p) \sim q^0$

- canonical field dimensions:

$$[\phi_c] = \frac{d-2}{2} < [\phi_q] = \frac{d+2}{2}$$

- action is dimensionless: phase e^{iS} in the functional integral
- $d > 2$: **couplings with more than two quantum fields irrelevant** in the RG sense
- massive simplification: semi-classical **Martin-Siggia-Rose-Janssen-de Dominicis action**

$$S = \int_{t,\mathbf{x}} \left\{ \underbrace{\phi_q^* \frac{\delta \bar{S}[\phi_c]}{\delta \phi_c^*}}_{\text{linear}} + c.c. + \underbrace{i2\gamma \phi_q^* \phi_q}_{\text{quadratic}} \right\} \quad \bar{S} = \int_{t,\mathbf{x}} \{ \underbrace{\phi_c^* i \partial_t \phi_c}_{\text{Hermitian}} - \underbrace{\mathcal{H}_c + i\mathcal{H}_d}_{\text{anti-Hermitian}} \}$$

Relation to driven-dissipative Gross-Pitaevskii-equation

- Equivalence of semiclassical theory and Langevin equations
 - structure of action in semiclassical limit (Martin-Siggia-Rose-Janssen-de Dominicis functional integral)

$$Z = \int \mathcal{D}[\phi_c, \phi_c^*, \phi_q, \phi_q^*] e^{iS[\phi_c, \phi_c^*, \phi_q, \phi_q^*]}$$

$$S = \int_{t,\mathbf{x}} \left\{ \phi_q^* \frac{\delta \bar{S}[\phi_c]}{\delta \phi_c^*} + c.c. + i2\gamma \phi_q^* \phi_q \right\}$$

$$\bar{S} = \int_{t,\mathbf{x}} \{ \phi_c^* i \partial_t \phi_c - \mathcal{H}_c + i \mathcal{H}_d \}$$

- Decouple the quantum field (undoing Gaussian integration)

$$Z = \int \mathcal{D}[\xi, \xi^*] e^{-\frac{1}{2\gamma} \int_{t,\mathbf{x}} \xi^* \xi} \int \mathcal{D}[\phi_c, \phi_c^*, \phi_q, \phi_q^*] e^{i \left[\phi_q^* \underbrace{\left(i \partial_t \phi_c - \frac{\delta \mathcal{H}_c}{\delta \phi_c^*} + i \frac{\delta \mathcal{H}_d}{\delta \phi_c^*} - \xi \right)}_{\frac{\delta \bar{S}}{\delta \phi_c^*}} + c.c. \right]}$$

- Recognize Fourier representation of delta-constraint: Langevin equation

$$Z = \int \mathcal{D}[\xi, \xi^*] e^{-\frac{1}{2\gamma} \int_{t,\mathbf{x}} \xi^* \xi} \int \mathcal{D}[\phi_c, \phi_c^*] \delta \left(i \partial_t \phi_c - \frac{\delta \mathcal{H}_c}{\delta \phi_c^*} + i \frac{\delta \mathcal{H}_d}{\delta \phi_c^*} - \xi \right) \delta(c.c.)$$

➔ noise averaging


➔ at each instant
of time:

➔ driven-dissipative Gross-Pitaevski equation

➔ cf. deterministic limit: noise added, system
explores many configurations


Semiclassical limit and exciton-polariton model

- example of “weak” universality (loss of memory of microscopic physics)





- ➔ many microscopic models collapse to an effective low energy model
- ➔ form dictated by microscopic symmetries
- ➔ longer wavelength behavior to be determined by calculation

Overview: Langevin equations, Lindblad equation, Keldysh integral




* effects of phase coherence still present (cf. BEC as classical wave)

“What is non-equilibrium about it?”



$$Z = \int \mathcal{D}(\Phi_+, \Phi_-) e^{i(S_M[\Phi_+, \Phi_-])}$$



“What is non-equilibrium about it?”

- different notions of ‘non-equilibrium’

Time evolution

- ➔ time translation invariance broken (e.g. thermalization, Floquet..) [see lectures by Anatoly and Norman!](#)

Stationary states (considered here)

- ➔ flux equilibria
 - **not** in static observables:

$$\rho = e^{-\beta H} / \text{tr} e^{-\beta H}$$


- ➔ any positive semidefinite Hermitian operator can be written like this
- dynamical observables, e.g.:

$$\langle \psi^\dagger(t) \psi(0) \rangle \quad \psi(t) = e^{iHt} \psi e^{-iHt}$$

- ➔ non-equilibrium conditions are encoded in the **generator of dynamics**
 - ➔ thermal equilibrium realized if generator of dynamics coincides with statistical weight
 - ➔ otherwise must expect non-equilibrium conditions (Lindbladian)

“What is non-equilibrium about it?”

- non-equilibrium stationary states:
 - **open system**: is it the coupling to a bath $\rightarrow$ irreversibility?
 - $\Rightarrow$ no, can be compatible with thermal equilibrium (Caldeira-Leggett Models)
 - **driven & open system**: coupling to multiple incompatible baths (e.g. different temperatures, chemical potential, driving strength...)



- result from independent microscopic processes
 - system is 'confused' which bath to thermalise to
 - some fluxes through the system
- $\Rightarrow$ driven open nature incompatible with thermal equilibrium
- $\Rightarrow$ how to sharply quantify?

“What is non-equilibrium about it?": Use symmetry!

- more formally: quantum master equation

$$\partial_t \rho = \underbrace{-i[H, \rho]}_{\Rightarrow S_H} + \underbrace{\mathcal{D}[\rho]}_{\Rightarrow S_{\mathcal{D}}}$$

- equivalent Keldysh functional integral: $Z = \int \mathcal{D}\phi_{\pm} e^{i(S_H[\phi_{\pm}] + S_{\mathcal{D}}[\phi_{\pm}])}$

- equilibrium dynamics microscopically generated by a **time-independent (undriven) Hamiltonian** alone

$$S_{\mathcal{D}} = 0$$

- ➔ **symmetry** of Keldysh action under discrete transformation

L. Sieberer, A Chiocchetta, U. Täuber, A. Gambassi, SD PRB (2015); F. Haehl, R. Loganayagam, M. Rangamani, JHEP (2016); M. Crossley, P. Glorioso, H. Liu, JHEP (2016)

$\mathcal{T}_{\beta} : \quad \phi_{\pm}(t, \mathbf{x}) \rightarrow \phi_{\pm}(-t + i\beta/2, \mathbf{x}), \quad i \rightarrow -i$

 $\mathcal{T}_{\beta}^2 = 1 \qquad \beta = 1/T$

- associated “Ward identities” are equilibrium quantum **fluctuation-dissipation relations** of arbitrary order



e.g. single-particle sector

$G^K(\omega, \mathbf{q}) = (2n_B(\omega/T) + 1)[G^R(\omega, \mathbf{q}) - G^A(\omega, \mathbf{q})]$

correlations Bose distribution responses

any order $\Leftrightarrow$ detailed balance
 $\Leftrightarrow$ global thermal equilibrium

- ➔ the Lindbladian ($S_{\mathcal{D}}$) violates this symmetry and therefore detailed balance **explicitly**
- ➔ intuition: no global bath to thermalise to

Equilibrium symmetry: Semiclassical limit

- Full symmetry:

$$\begin{aligned}\mathcal{T}_\beta : \quad \phi_\pm(t, \mathbf{x}) &\rightarrow \phi_\pm(-t + i\beta/2, \mathbf{x}), \quad i \rightarrow -i & \beta = 1/T \\ &= e^{\pm i\frac{\beta}{2}\partial_t} \phi_\pm(-t, \mathbf{x})\end{aligned}$$

- semiclassical limit: T large $\Rightarrow$ $e^{\pm i\frac{\beta}{2}\partial_t} \approx 1 \pm i\frac{\beta}{2}\partial_t$

- action on the fields:

irrelevant by power counting

$$\mathcal{T}_\beta \phi_c(t, \mathbf{x}) = \phi_c^*(-t, \mathbf{x}) + \frac{i}{2T} \partial_t \phi_q^*(-t, \mathbf{x}),$$

reproduces classical result

$$\mathcal{T}_\beta \phi_q(t, \mathbf{x}) = \phi_q^*(-t, \mathbf{x}) + \frac{i}{2T} \partial_t \phi_c^*(-t, \mathbf{x})$$

H. K. Janssen (1976); C. Aron
et al, J Stat. Mech (2011)


➡ obtain **geometric interpretation** of the equilibrium symmetry

Geometric interpretation: equilibrium vs. non-equilibrium dynamics

- in semi-classical **Martin-Siggia-Rose-Janssen-de Dominicis** action

$$S = \int_{t,\mathbf{x}} \left\{ \phi_q^* \frac{\delta \bar{S}[\phi_c]}{\delta \phi_c^*} + c.c. + i2\gamma \phi_q^* \phi_q \right\} \quad \bar{S} = \int_{t,\mathbf{x}} \{ \phi_c^* i \partial_t \phi_c - \mathcal{H}_c + i \mathcal{H}_d \}$$

$$\mathcal{H}_\alpha = \int d^3x [r_\alpha |\phi|^2 + K_\alpha |\nabla \phi|^2 + \lambda_\alpha |\phi|^4] \quad \alpha = c, d$$




example: two-body processes λ

$\text{Re}\lambda$ elastic two-body collisions

$\text{Im}\lambda$ inelastic two-body losses


Geometric interpretation: equilibrium vs. non-equilibrium dynamics

equilibrium dynamics



- coherent and dissipative dynamics may occur simultaneously
- but they are not independent


non-equilibrium dynamics



- coherent and dissipative dynamics do occur simultaneously
- they result from different dynamical resources

➔ what are the physical consequences of the spread in the complex plane?

Application: Fate of BKT physics in driven open quantum systems



E. Altman, L. Sieberer, L. Chen, SD, J. Toner, PRX (2015)

G. Wachtel, L. Sieberer, SD, E. Altman, PRB (2016)

L. Sieberer, G. Wachtel, E. Altman, SD, PRB (2016)

L. He, L. Sieberer, E. Altman, SD, PRB (2015)

L. He, L. Sieberer, SD, PRL (2017)

Microscopic
Quantum Optics

“Thermodynamic”
Many-body physics

Long wavelength
Statistical mechanics

Phase transitions in two dimensions

- continuous symmetry U(1): no spontaneous symmetry breaking, but a phase transition



- correlations

$$\langle \phi^*(\mathbf{x}) \phi(0) \rangle \sim r^{-\frac{1}{2\pi K}}$$

$$\sim e^{-r/\xi}$$





- responses: superfluidity

$$\rho_s \neq 0$$

$$\rho_s = 0$$

- BKT (Berezinskii-Kosterlitz-Thouless) transition: unbinding of vortex-antivortex pairs


J. M. Kosterlitz, D. J. Thouless J. Phys. C (1973)



matter wave interferometry:
Z. Hadzibabic et al. Nature (2006)

... fate in driven open condensates?

Short reminder: Algebraic correlations



- correlations

$$\langle \phi^*(\mathbf{x}) \phi(0) \rangle \sim r^{-\frac{1}{2\pi K}} \quad \sim e^{-r/\xi}$$


- physical reason: gapless spin wave fluctuations

- spin wave action $S = \frac{K}{2} \int d^2x (\nabla \theta)^2 = \frac{K}{2} \int \frac{d^2q}{(2\pi)^2} q^2 \theta(-\mathbf{q}) \theta(\mathbf{q})$

- phase-amplitude decomposition $\phi(\mathbf{x}) = \rho(\mathbf{x})^{1/2} e^{i\theta(\mathbf{x})} \approx \sqrt{n_0} e^{i\theta(\mathbf{x})}$

$$\langle \phi^*(\mathbf{x}) \phi(0) \rangle \approx n_0 \langle e^{i(\theta(\mathbf{x}) - \theta(0))} \rangle = n_0 e^{-\frac{1}{2} \langle (\theta(\mathbf{x}) - \theta(0))^2 \rangle}$$


- phase correlator $\frac{1}{2} \langle (\theta(\mathbf{x}) - \theta(0))^2 \rangle = \int^{1/a} \frac{d^2q}{(2\pi)^2} \frac{(e^{iqr} - 1)}{Kq^2} = \frac{1}{2\pi K} \log(r/a)$



Description: Effective model

- mesoscopic starting point: driven-dissipative stochastic Gross-Pitaevski equation

$$i\partial_t\phi = \left[-\frac{\nabla^2}{2m} - \mu + i(\gamma_p - \gamma_l) + (g - i\kappa)|\phi|^2 \right] \phi + \zeta$$



$$\phi = \rho e^{i\theta}$$

- effective low frequency dynamics see also: G. Grinstein et al., PRL (1993)

$$\partial_t\theta = D\nabla^2\theta + \lambda(\nabla\theta)^2 + \xi$$

phase diffusion

phase nonlinearity

Markov noise

form of the KPZ equation

Kardar, Parisi, Zhang, PRL (1986)

- meaning: non-linear spin wave mode
- nonlinearity: single-parameter measure of non-equilibrium strength (ruled out in equilibrium by symmetry)


microphysics

quantum master equation

stochastic GPE

KPZ equation

macrophysics



KPZ equation: A paradigm of non-equilibrium stat mech


- originally: describes stochastic roughening of surface height $h(\mathbf{x}, t)$

$$\partial_t h = D \nabla^2 h + \lambda (\nabla h)^2 + \xi$$

smoothens nonlinear growth noise

Kardar, Parisi, Zhang, PRL (1986)
Review: Krug, Adv. Phys. (1997)

- simplest physical scenario



KPZ equation: A paradigm of non-equilibrium stat mech


- originally: describes stochastic roughening of surface height $h(\mathbf{x}, t)$

$$\partial_t h = D \nabla^2 h + \lambda (\nabla h)^2 + \xi$$

smoothens nonlinear growth noise

Kardar, Parisi, Zhang, PRL (1986)
Review: Krug, Adv. Phys. (1997)

- multiple physical contexts



defect growth in liquid
crystals

drive: electric field

from Takeuchi et al.,
Scientific Reports (2011)



bacterial colony growth

drive: sugar

Wakita et al., J. Phys. Jpn.
Soc. (1997)





burning paper

drive: oxygen


Maunuksela et al., PRL
(1997)

Physical implication I: Smooth KPZ fluctuations

- How important are non-equilibrium conditions at large distance?
- behavior of non-equilibrium strength under coarse graining (RG flow)




- general trend: non-equilibrium effects in systems with gapless mode are
 - enhanced in $d = 1, 2$
 - softened in $d = 3$ (below a threshold)



Physical implication I: Smooth KPZ fluctuations


- How important are non-equilibrium conditions at large distance?
- behavior of non-equilibrium strength under coarse graining (RG flow)



- 2D: implication: a length scale is generated


$$L_* = a_0 e^{\frac{16\pi D^3}{\lambda^2 \Delta}}$$

microscopic (healing)
length




2D: Absence of algebraic order out of equilibrium

- generated length scale distinguishes two regimes: $L_* = a_0 e^{\frac{16\pi D^3}{\lambda^2 \Delta}}$



- algebraic order **absent** in any two-dimensional driven open system at the largest distances
- but crossover scale **exponentially large** for small deviations from equilibrium
- observation in 1D systems (KPZ scaling exponents found)



$\beta = 1/3$ temporal scaling

Q. Fontaine et al., Nature (Aug 24 2022)


spatial scaling $\chi = 1/2$

Physical implications II: Non-equilibrium Kosterlitz-Thouless


- KPZ equation for **phase variable**

$$\partial_t \theta = D \nabla^2 \theta + \lambda (\nabla \theta)^2 + \xi$$

- compact nature of phase allows for vortex defects in 2D!
- key ingredient of Kosterlitz-Thouless transition




vortex




anti-vortex

$$F = E - TS$$

low T:
(binding) energy dominates



high T:
entropy dominates





→ how is this scenario modified in the driven system?



Mini-review: BKT transition

low temperature

high temperature



- BKT transition: unbinding of vortex-antivortex pairs
J. M. Kosterlitz, D. J. Thouless J. Phys. C (1973)



- Single vortex picture: balance of energy (deterministic) and entropy (statistic)
 - Low T: vortices and antivortices bound in neutral pairs (irrelevant at long distance)
 - Q: when is it favorable (free energy minimum) to have **unbound** vortices?

- energy of single free vortex:

vortex configuration: mapping $(r, \varphi) \rightarrow \theta(r, \varphi) = \varphi \implies \nabla\theta = \frac{1}{r}\hat{e}_\phi\partial_\varphi\theta(r, \varphi) = \frac{1}{r}\hat{e}_\phi$

$$\implies E = \frac{K}{2} \int d^2x (\nabla\theta)^2 = \pi K \int_a^L dr r \frac{1}{r^2} = \pi K \log(L/a)$$

- entropy: sum all equally probable possibilities of placing vortices in 2D plane at minimal distance a:

$$S = -k_B \sum_i p_i \log p_i = k_B \log(L/a)^2$$

- free energy $F = E - TS = (\pi K - 2k_B T) \log(L/a)$

- vortex proliferation above KT critical temperature $T_{KT} = \frac{\pi K}{2k_B}$

→ but out of equilibrium: no free energy at hand!

→ field theory approach (analogous Kosterlitz' real space RG for vortices)

J. M. Kosterlitz, J. Phys. C (1974)

Compact KPZ

L. Sieberer, G. Wachtel, E. Altman, SD, PRB (2016)
G. Wachtel, L. Sieberer, SD, E. Altman, PRB (2016)

- wait a second — we ignored a fundamental symmetry of polaritons so far: local discrete gauge invariance

$$\phi(t, \mathbf{x}) = \rho(t, \mathbf{x}) e^{i\theta(t, \mathbf{x})} \quad \theta_{t, \mathbf{x}} \mapsto \theta_{t, \mathbf{x}} + 2\pi n_{t, \mathbf{x}}$$

- Teaching symmetry to KPZ equation:

$$\theta_{t+\epsilon, \mathbf{x}} = \theta_{t, \mathbf{x}} + \epsilon (\mathcal{L}[\theta]_{t, \mathbf{x}} + \eta_{t, \mathbf{x}}) + 2\pi n_{t, \mathbf{x}}$$

lattice regularized deterministic term

stochastic **difference**
equation

Non-equilibrium
electrodynamic duality:

Electric field $\Leftrightarrow$ smooth phase fluct. (KPZ)

Charges $\Leftrightarrow$ vortices

Dynamical & non-equilibrium analog of
Kosterlitz-Thouless construction



$$Z = \sum_{\{\tilde{n}_{t, \mathbf{x}}\}} \int \mathcal{D}[\theta] e^{iS[\theta, \tilde{n}]}$$

discrete noise MSRJD
functional integral

non-eq. lattice gauge theory



$$Z \propto \sum_{\{n_{vX}, \tilde{n}_{vX}, \mathbf{J}_{vX}, \tilde{\mathbf{J}}_{vX}\}} \int \mathcal{D}[\phi, \tilde{\phi}, \mathbf{A}, \tilde{\mathbf{A}}] e^{iS[\phi, \tilde{\phi}, \mathbf{A}, \tilde{\mathbf{A}}, n_v, \tilde{n}_v, \mathbf{J}_v, \tilde{\mathbf{J}}_v]}$$


non-equilibrium
electrodynamic theory

➡ Q: What is the effective theory for vortices out of equilibrium?

Effective theory for a single vortex-antivortex pair

- equation of motion for a single vortex-antivortex pair

$$\frac{d\mathbf{r}}{dt} = -\mu \nabla V(r) + \boldsymbol{\xi}$$



equilibrium: Coulomb potential (2D)

$$V(\mathbf{r}) = \frac{1}{\varepsilon} \ln(r/a)$$

$\varepsilon \equiv K$
(sorry..)

length scale:

$$L_v = a_0 e^{\frac{2D}{\lambda}}$$

see also: I Aranson
et al., PRB (1998)
two-vortex problem

$$V(\mathbf{r}) \approx \frac{1}{\varepsilon} \ln(r/a) - \frac{\lambda^2}{12\varepsilon^3 D^2} (\ln(r/a))^3$$


driven-dissipative system

➔ noise-activated unbinding for a single pair (at exp small rate)

Many pairs: Corrections to Kosterlitz-Thouless flow


$$\frac{dK}{d\ell} = \frac{2\pi^2 y^2}{T} \quad \frac{dy}{d\ell} = \left[2 - \frac{1}{2KT} + \frac{\lambda^2}{4K^2 T^2} \left(\frac{1}{4} + \ell \right) \right] y \quad \frac{dT}{d\ell} = \frac{\lambda^2 T}{2K^2 T^2} \left(\frac{1}{4} + \ell \right)$$

- parameter y : $\sim$ probability to create single vortex at distance l (flow parameter)



Many pairs: Corrections to Kosterlitz-Thouless flow

$$\frac{dK}{d\ell} = \frac{2\pi^2 y^2}{T} \quad \frac{dy}{d\ell} = \left[2 - \frac{1}{2KT} + \frac{\lambda^2}{4K^2 D^2} \left(\frac{1}{4} + \ell \right) \right] y \quad \frac{dT}{d\ell} = \frac{\lambda^2 T}{2K^2 D^2} \left(\frac{1}{4} + \ell \right)$$



Competing length scales and suppression of KT


- two emergent length scales in complementary approaches:

$$L_* = a_0 e^{\frac{16\pi D^3}{\lambda^2 \Delta}}$$

KPZ length

$$L_v = a_0 e^{\frac{2D}{\lambda}}$$

vortex length




- numerical confirmation of two-scale scenario in 1D (defects: vortices in (1+1)D space-time)

L. He, L. Sieberer, SD PRL (2017)

- 2D simulations demanding

Kosterlitz-Thouless physics fragile to non-equilibrium perturbation

A phase transition driven by non-equilibrium strength




L. He, L. Sieberer, E. Altman, SD, PRB (2015)


L. He, L. Sieberer, SD, PRL (2017)

Sequence of Scales

- direct numerical solution of driven-dissipative GPE in one dimension
- Study temporal instead of spatial coherence function






numerical evidence



➔ no spatial vortices in 1D ➔ what causes the emergent length scale beyond KPZ?

Space-time vortices in 1D

- Physical origin: compactness of phase field



topologically nontrivial phase field configurations on (1+1)D space-time plane

- unbound** at infinitesimal noise level (weak non-equilibrium)


- interaction potential: $(\partial_t + D\partial_x^2)^{-1} \sim (Dt)^{-1/2} e^{-x^2/(4Dt)}$ cf. 2D static equilibrium: $\nabla^{-2} \sim \log(|\mathbf{x}|)$

- explains qualitative features

- temporal scaling** (random uncorrelated charges) $\langle \psi^*(x, t') \psi(x, t) \rangle \sim e^{-c|t-t'|}$
- noise level dependence** of crossover scale $T_v \sim e^{E_c/\sigma}$ (mapping to static 2D active smectic A liquid crystal)

Strong non-equilibrium: Compact KPZ vortex turbulence



- In search of the phase diagram for XP condensates




Strong non-equilibrium: Compact KPZ vortex turbulence

- In search of the phase diagram for XP condensates: 1+1 dimensions

L. He, L. Sieberer, SD PRL (2017)



→ deterministic limit: how does the system generate its own noise?



Phase transition driven by non-equilibrium strength

- qualitative reason: competition in lattice regularized model (compact KPZ)

$$\partial_t \vec{\chi} = \underbrace{S \vec{V}_E[\vec{\chi}]}_{\text{equilibrium}} + \underbrace{A \vec{V}_N[\vec{\chi}]}_{\text{non-equilibrium}} + \vec{\xi}$$

lattice site

$$\chi_i = \nabla \theta_i = \theta_{i+1} - \theta_i$$

$$V_{E/N,i} = \frac{\delta}{\delta \chi_i} \mathcal{V}_{E/N}$$


$$\mathcal{V}_E = K \sum_j \cos \chi_j \quad \mathcal{V}_N = \lambda \sum_j \sin \chi_j$$

$$D_{ij} = \delta_{ij+1} - \delta_{ij}$$


$$S = D + D^T = S^T$$

$$A = D - D^T = -A^T$$


fixed point
structures




attractive



repulsive



hyperbolic



elliptic

=> emergent Liouville thm.


$$\text{div}[A \vec{V}_N] = 0$$

- high dimensional configuration space
- transition to chaos

Further instances: Universality in driven open quantum systems


- criticality in non-equilibrium ϕ^4 model Sieberer et al., PRL (2013)

→ dynamical fine-structure distinguishing eq. from non-eq.




- Markovian quantum criticality Marino, SD, PRL (2016)

→ new fixed point in dark state models with quantum scaling




- phase transitions in open Floquet systems Mathey, SD, PRL (2019)

→ absence of criticality (dual to Kibble-Zurek mechanism)



- coupled Ising models Young et al., PRX (2020)


→ new fixed point for strong non-equilibrium drive



- phase transitions with exceptional points Fruchart et al., Nature (2021)


→ fluctuation induced first order transition

Zelle, Daviet, Rosch, SD arxiv (2023)



Summary lecture I & II

Review: L. Sieberer, M. Buchhold, SD, *Keldysh Field Theory for Driven Open Quantum Systems*, Reports on Progress in Physics (2016)




Many-Body Master Equation

1-1
mapping

Keldysh functional
integral

- mapping opens up QFT toolbox:
 - symmetries: eq. vs. non-eq.
 - control of IR fluctuations: understanding low dimensional gapless phases out of equilibrium
 - flexible choice of degrees of freedom: KPZ vs. vortices
- next lecture: open systems $\rightarrow$ measurements in many body systems
 - weak continuous measurements
 - replica approach and replica Keldysh functional integral
 - BKT transition of replica fluctuations for monitored fermions





School on quantum many-body
phenomena out of equilibrium
August 25/26, 2023
ICTP Trieste, Italy



Lecture III

Lindblad-Keldysh 2.0: Measurement Induced Phase of Fermions


Sebastian Diehl

Institute for Theoretical Physics, University of Cologne

Introduction

Small quantum systems: Measurements

- two types of quantum dynamics





for measurement observable $\hat{M} = \sum_{\lambda} m_{\lambda} |\lambda\rangle \langle \lambda| \equiv \sum_{\lambda} m_{\lambda} \hat{P}_{\lambda}$

- dynamics non-trivial (eigenstates not shared) once $[\hat{H}, \hat{M}] \neq 0$

Many-body systems: Phase transitions

- non-commuting operators lead to (quantum) phase transitions

$$\hat{H} = \hat{H}_1 + g\hat{H}_2 \quad [\hat{H}_1, \hat{H}_2] \neq 0$$



e.g. Mott-insulator to superfluid transition in cold atoms


→ combine measurement and many particles: similar scenario?

Entanglement Phase Transitions in Random Circuits


- model and key ingredients:

- randomly chosen local entangling unitary gates
- projective local measurement of non-commuting observables


Skinner, Ruhman, Nahum
PRX (2019)



Li, Chen, Fisher, PRB
(2018, 2019)



- basic picture: competition in many-body context (single trajectory, exactly local meas. of σ_i^z)



- Procedure

- track single quantum trajectories (pure states)
- compute the quantity of interest (e.g. entanglement entropy)
- average over trajectory ensemble

non-commuting


$$|\{\sigma_i\}\rangle\langle\{\sigma_i\}| \sim \mathbf{1}$$

Entanglement Phase Transitions in Random Circuits


- model and key ingredients:

- randomly chosen local entangling unitary gates
- projective local measurement of non-commuting observables


Skinner, Ruhman, Nahum
PRX (2019)



Li, Chen, Fisher, PRB
(2018, 2019)



- basic picture: competition in many-body context (single trajectory, exactly local meas. of σ_i^z)



- Procedure


- track single quantum trajectories (pure states)
- compute the quantity of interest (e.g. entanglement entropy)
- average over trajectory ensemble

non-commuting

$$|\{\sigma_i\}\rangle \langle \{\sigma_i\}| \sim \mathbf{1}$$

$\Rightarrow$ Phase transition in entanglement growth at **finite** competition ratio g

Outline Lecture III



- measurements: description?

- strong projective vs. weak continuous measurements
- ‘observables’

- many-body problem: phase transitions?

- BKT type phase transition from critical to area law phase
- signatures beyond entanglement entropy

- how to understand?

- replicated Lindblad equation
- replicated Keldysh field theory


- how to observe?

- post vs. preselection

➡ Quantum phase transition in trajectory wavefunction, revealed in non-linear-in-state observables

Theory Background:


Strong projective vs. weak continuous measurements



Projective vs. Weak Measurements

Review: Jacobs, Steck, Contemp. Phys. (2006)

- projective measurement: acquire full knowledge about observable



- example: projective position measurement

- measurement observable

$$\hat{X} = \int dx x |x\rangle \langle x| \equiv \int dx x \hat{P}(x)$$

- measurement outcome $\lambda = x_0$

- projector onto that state


$$\hat{P}(x_0) = |x_0\rangle \langle x_0| = \int dx \delta(x - x_0) |x\rangle \langle x|$$

➡ know post-measurement position x_0 with certainty

Projective vs. Weak Measurements

Review: Jacobs, Steck, Contemp. Phys. (2006)

- projective measurement: acquire full knowledge about observable



→ know post-measurement position x_0 with certainty

- instantaneous collapse of wave function, but real measurements take time: **continuous observation**


$$\hat{P}(x_0) \rightarrow \hat{A}(x_0) = \left(\frac{2\gamma\Delta t}{\pi}\right)^{1/4} \int dx e^{-\gamma\Delta t(x-x_0)^2} |x\rangle\langle x| = \left(\frac{2\gamma\Delta t}{\pi}\right)^{1/4} e^{-\gamma\Delta t(\hat{x}-x_0)^2}$$

→ **strong projective measurement** obtains for $\Delta t \rightarrow \infty$

→ **weak continuous measurement** obtains for small Δt , but large number of repetitions N :

$$\Delta t \rightarrow 0, \quad N \rightarrow \infty, \quad \Delta t \cdot N \rightarrow \text{const.}$$

cf. functional integral construction!



Probabilistic Character of Weak Measurements

Review: Jacobs, Steck, Contemp. Phys. (2006)

- probability to measure x_0 on state $|\psi\rangle = \int dx \psi(x) |x\rangle$ in time $\Delta t \rightarrow 0$

$$p(x_0) = ||\hat{A}(x_0)|\psi\rangle|| = \text{tr}[\hat{A}^\dagger(x_0)\hat{A}(x_0)|\psi\rangle\langle\psi|] \quad \hat{A}(x_0) = \left(\frac{2\gamma\Delta t}{\pi}\right)^{1/4} e^{-\gamma\Delta t(\hat{x}-x_0)^2}$$

- the measurement outcome x_0 is a Gaussian random variable with

- expectation value $\langle\langle x_0 \rangle\rangle \equiv \int dx_0 x_0 p(x_0) = \left(\frac{2\gamma\Delta t}{\pi}\right)^{1/2} \int dx_0 \int dx x_0 e^{-2\gamma\Delta t(x-x_0)^2} |\psi(x)|^2 = \int dx x |\psi(x)|^2 = \langle\hat{x}\rangle$

- variance $\langle\langle x_0^2 \rangle\rangle - \langle\langle x_0 \rangle\rangle^2 = \frac{1}{4\gamma\Delta t}$

quantum mechanical expectation

- stochastic formulation (cf. Fokker-Planck vs. Langevin): parameterize


$$x_0 = \langle\hat{x}\rangle + \frac{\Delta W}{2\gamma\Delta t} \quad \text{Gaussian random variable} \quad \begin{aligned} \overline{\Delta W} &= 0 \\ \overline{\Delta W \Delta W} &= \gamma\Delta t \end{aligned} \quad \implies \Delta W \sim \sqrt{\Delta t}$$

- stochastic update of wave function: $|\psi_{t+\Delta t}\rangle = \frac{\hat{A}(x_0)|\psi_t\rangle}{||\hat{A}(x_0)|\psi_t\rangle||}$

- for short observation times: expand $|\psi_{t+\Delta t}\rangle = \hat{A}(x_0)|\psi_t\rangle$ to linear order $\Delta t \equiv dt \rightarrow 0$

$$d|\psi_t\rangle \equiv |\psi_{t+dt}\rangle - |\psi_t\rangle = \left\{ \left[-\frac{1}{2}\gamma(\hat{x} - \langle\hat{x}\rangle)^2\right]dt + (\hat{x} - \langle\hat{x}\rangle)dW \right\} |\psi\rangle$$

Weak Measurements: Stochastic Schrödinger Equation




- governed by **stochastic Schrödinger equation** for **quantum trajectory** $|\psi_t\rangle$ V. P. Belavkin (1987)

$$d|\psi_t\rangle = dt(-i\hat{H} - \frac{\gamma}{2} \sum_l (\hat{n}_l - \langle \hat{n}_l \rangle_t)^2 |\psi_t\rangle) + \sum_l \underbrace{dW_l (\hat{n}_l - \langle \hat{n}_l \rangle_t)}_{\text{multiplicative noise}} |\psi_t\rangle$$


$\langle \hat{n}_l \rangle_t = \langle \psi_t | \hat{n}_l | \psi_t \rangle$

- Hamiltonian added, many degrees of freedom, $\hat{n}_l = \hat{n}_l^\dagger$
 - works for measurement operators with discrete spectrum (measurement record continuous, e.g. Stern-Gerlach)
 - measurement only dynamics $H=0$: **measurement dark states**
 - dW ‘multiplicative noise’: inactive when $\hat{n}_l |\psi_t\rangle = \langle \hat{n}_l \rangle_t |\psi_t\rangle$
 - e.g. in eigenstate $\hat{n}_l |\psi_t\rangle = n_l |\psi_t\rangle$

dark state of measurement operator
- **continuous collapse**: convergence to measurement eigenstate for long times (more gen. for) $[\hat{H}, \hat{n}_l] = 0$




Weak Measurements: Stochastic Schrödinger Equation



- governed by **stochastic Schrödinger equation** for **quantum trajectory** $|\psi_t\rangle$ [Belavkin \(1987\); Gisin, Percival \(1993\)](#)

$$d|\psi_t\rangle = dt(-i\hat{H} - \frac{\gamma}{2} \sum_l (\hat{n}_l - \langle \hat{n}_l \rangle_t)^2 |\psi_t\rangle + \sum_l \underbrace{dW_l (\hat{n}_l - \langle \hat{n}_l \rangle_t)}_{\langle \hat{n}_l \rangle_t = \langle \psi_t | \hat{n}_l | \psi_t \rangle} |\psi_t\rangle$$

- Hamiltonian added, many degree of freedom, $\hat{n}_l = \hat{n}_l^\dagger$
- works for measurement operators with discrete spectrum (measurement record continuous, e.g. Stern-Gerlach)
- Can we expect a measurement induced phase transition similar to projective measurements? Yes!
 - no competition => no phase transition $\hat{A}(x_0) = (\frac{2\gamma\Delta t}{\pi})^{1/4} e^{-\gamma\Delta t(\hat{x}-x_0)^2} \xrightarrow{\Delta t \rightarrow \infty} |x_0\rangle\langle x_0|$
 - Demonstrated numerically [Szyniszewski, Romito, Schomerus, PRB \(2019\)](#)



- expect **no phase transition** upon taking temporal continuum limit
- continuum limit useful for analytical approach measurement induced phase transitions (e.g. Keldysh field theory approach)

Monitored Dynamics: Extracting Information

- $|\psi_t\rangle$ is a **random variable** -> needs statistical analysis
- **stochastic Schrödinger equation** for projector $\hat{\rho}_t = |\psi_t\rangle\langle\psi_t|$

$$d\rho_t = \underbrace{dt(-i[\hat{H}, \rho_t] + \frac{\gamma}{2} \sum_l (2\hat{n}_l \rho_t \hat{n}_l - \{\hat{n}_l^2, \hat{\rho}_t\}))}_{\text{deterministic part}} + \sum_l dW_l \{\hat{n}_l - \langle\hat{n}_l\rangle_t, \rho_t\}$$



→ statistical analysis: consider trajectory ensemble

- usual observables:

$$\begin{array}{ccc} \langle\hat{O}\rangle = \langle\psi_t|\hat{O}|\psi_t\rangle = \text{tr}[\hat{O}\hat{\rho}_t] & \xrightarrow[\text{e.g. trajectories}]{\text{statistical average}} & \overline{\langle\hat{O}\rangle} = \overline{\langle\psi_t|\hat{O}|\psi_t\rangle} = \text{tr}[\hat{O}\overline{\hat{\rho}_t}] \\ \text{quantum average} & & \end{array}$$

- Problem: Hermitian measurement operators => $\overline{\hat{\rho}_t} \sim \mathbf{1}$

→ one way to see this: upon statistical average of equation of motion,

- deterministic, linear **Lindblad equation** for $\rho_t = \overline{|\psi_t\rangle\langle\psi_t|}$
- $\hat{n}_l$ Hermitian: heating to **infinite temperature** $\longrightarrow \rho_t \sim \mathbf{1}$

Monitored Dynamics: Extracting Information

- $|\psi_t\rangle$ is a **random variable** -> needs statistical analysis
- **stochastic Schrödinger equation** for projector $\hat{\rho}_t = |\psi_t\rangle\langle\psi_t|$

$$d\rho_t = dt(-i[\hat{H}, \rho_t] + \frac{\gamma}{2} \sum_l (2\hat{n}_l \rho_t \hat{n}_l - \{\hat{n}_l^2, \hat{\rho}_t\}) + \sum_l dW_l \{\hat{n}_l - \langle\hat{n}_l\rangle_t, \rho_t\})$$

→ statistical analysis: consider trajectory ensemble

- usual observables:

$$\begin{array}{ccc} \langle\hat{O}\rangle = \langle\psi_t|\hat{O}|\psi_t\rangle = \text{tr}[\hat{O}\hat{\rho}_t] & \xrightarrow[\text{e.g. trajectories}]{\text{statistical average}} & \overline{\langle\hat{O}\rangle} = \overline{\langle\psi_t|\hat{O}|\psi_t\rangle} = \text{tr}[\hat{O}\overline{\hat{\rho}_t}] \\ \text{quantum average} & & \end{array}$$

- Problem: Hermitian measurement operators => $\overline{\hat{\rho}_t} \sim \mathbf{1}$

→ solution: use **state-dependent ‘observables’** $\overline{\langle\hat{O}(|\psi\rangle)\rangle} = \text{tr}\hat{O}(\overline{\hat{\rho}})\overline{\hat{\rho}}$

more promising, because in general $F(\overline{\hat{\rho}}) \neq \overline{F[\hat{\rho}]}$


- examples:

- von Neumann entropy

$$\overline{S_{vN}(l, L)} = \overline{\langle\log(\rho_A)\rangle}$$

- correlation function

$$\overline{\langle\hat{n}_i\rangle\langle\hat{n}_j\rangle}$$




arbitrarily high power of state projector

quadratic in state projector

An example: TLS / one fermion on two sites

- toy model: trajectory evolution of single fermion on two sites

$$|\psi_{t+dt}\rangle = |\psi_t\rangle - i dt \hat{H}_{\text{eff}} |\psi_t\rangle + \sum_{l=1}^2 dW_l (\hat{n}_l - \langle \hat{n}_l \rangle_t) |\psi_t\rangle$$




$$\hat{H}_{\text{eff}} = \hat{H} - i\hat{K} \quad \hat{H} = -J (c_1^\dagger c_2 + h.c.) \quad \hat{K} = \frac{\gamma}{2} \sum_{l=1}^2 (\hat{n}_l - \langle \hat{n}_l \rangle_t)^2$$

guiding physical picture:

- thermodynamic limit: pinning quantum phase transition may happen at sharply defined point
- signalled in nonlinear-in-state ‘observable’, like the covariance matrix


→ **pinning** to measurement eigenstate

- invisible in linear averages




→ vanishing time spent in eigenstate

- seen in **averaged trajectory covariance matrix**




Entanglement Phase Transition in a Monitored Fermion Chain

Hamiltonian:



entanglement growth


Monitoring:



entanglement saturation

O. Alberton, M. Buchhold, SD, PRL (2021)

Monitored Fermion Dynamics



- Weak continuous measurements in many-body system

$$d|\psi_t\rangle = dt(-i\hat{H} - \frac{\gamma}{2} \sum_l \hat{M}_l^2 |\psi_t\rangle + \sum_l dW_l \hat{M}_l |\psi_t\rangle)$$

Gaussian white noise g

- competition: $g = \frac{\gamma}{J}$



- unitary dynamics: hopping

- measurement operators $\hat{M}_l = \hat{n}_l - \langle \hat{n}_l \rangle_t$

- $H = 0$: evolution stops after collapse into **dark state**

$$H = -J \sum_l (c_l^\dagger c_{l+1} + c_{l+1}^\dagger c_l)$$

$$\hat{M}_l |\psi_t\rangle = 0 \quad \text{for} \quad \hat{n}_l |\psi_t\rangle = n_l |\psi_t\rangle$$


eigenstate of measurement operator

- volume law entanglement entropy

- area law entanglement entropy

$$S_{vN}(L/2, L) = \text{tr} \rho_A \log(\rho_A) \stackrel{t \rightarrow \infty}{\sim} L$$


$$S_{vN}(L/2, L) = s_0$$




- caveat: $|\psi_t\rangle$ is a random variable
- binary measurement outcomes generate extensive **configurational entropy**

- ‘observables’: entanglement entropy, traj. averaged correlators

Trajectory Ensemble Phase Diagram: Entanglement Entropy



Characterizing the Weak Monitoring Phase




- extended criticality: Connected correlation function

- emergent conformality: Mutual information


$$C_{i,i+l} = \langle \hat{n}_i \rangle \langle \hat{n}_{i+l} \rangle - \langle \hat{n}_i \hat{n}_{i+l} \rangle$$

$$\sim l^{-2}$$




$$\mathcal{I}(A, B) = \overline{S_{vN}(A)} + \overline{S_{vN}(B)}$$

$$+ \overline{S_{vN}(A \cup B)}$$



- captures all distinct phases:


$$C_{i,i+l} \sim \begin{cases} 0 & \text{for } H = 0 \\ \exp(-l/\xi) & \text{for } \gamma \gg J \\ l^{-2} & \text{for } \gamma \ll J \\ l^{-1} & \text{for } \gamma = 0 \end{cases}$$



conformally invariant critical point: Nahum et al. PRX (2019); Li Chen Fisher PRB (2019); Jian et al. PRB (2020);


➡ emergent conformally invariant critical phase for weak monitoring

Characterizing the Phase Transition



- effective central charge $c(\gamma)$


$$\overline{S_{vN}(l, L)} = \frac{c(\gamma)}{3} \log_2 \left[\frac{L}{\pi} \sin \left(\frac{\pi l}{L} \right) \right] + s(\gamma)$$



parameter dependent c

random systems: Cardy Jacobsen PRL (1997);
Refael, Moore PRL (2004)

- essential scaling of the central charge: scaling collapse above phase transition




➔ sudden jump reminiscent of BKT in 1+1 dimensions (Berezinskii-Kosterlitz-Thouless)

➔ BKT universal behavior

➔ numerically establishes **BKT type phase transition**

Replica Field Theory Approach to Measurement Induced Phase Transitions



M. Buchhold, Y. Minoguchi, A. Altland, SD, PRX 11, 041004 (2021)


microphysics






macrophysics

Continuum (1+1) dimensional Model

- approach problem from the weakly measured side



- model: phenomenologically motivated continuum limit for lattice fermions (weak measurement regime)




- weak measurement: Hamiltonian dominates, measurement as perturbation
- stack of almost decoupled patches (rotating wave approximation in real space and time)
- focus on patch with strongest backscattering $k_n = \pi/2$: **Dirac model**

Continuum (1+1) dimensional Model

- model: phenomenologically motivated continuum limit for lattice fermions (weak measurement regime)

- Hamiltonian: massless Dirac fermions $\hat{\Psi}_x = (\hat{\psi}_{R,x}, \hat{\psi}_{L,x})^T$



$$\hat{H} = i v \int_x \hat{\Psi}_x^\dagger \sigma_z \partial_x \hat{\Psi}_x$$

fermionic variant

bosonization
→
dictionary

$$\hat{H} = \frac{v}{2\pi} \int_x [(\partial_x \hat{\theta}_x)^2 + (\partial_x \hat{\phi}_x)^2]$$

phase density

bosonized variant

- measurement operators: local particle density $\rightarrow$ current and vertex operators

$$\text{rate } \gamma_1 : \quad \hat{O}_{1,x} = \Psi_x^\dagger \Psi_x = \hat{J}_x^{(0)}$$

$$\hat{O}_{1,x} = -\frac{1}{\pi} \partial_x \hat{\phi}_x \quad \text{linear gapless}$$

rate γ_2 : $\hat{O}_{2,x} = \Psi_x^\dagger \sigma_x \Psi_x$

$$\hat{O}_{2,x} = \underbrace{m \cos(2\hat{\phi}_x)}_{\mathcal{O}(1)} \quad \text{nonlinear}$$


common eigenstates: $\hat{\phi}_x |\Psi_D\rangle = \phi_x |\Psi_D\rangle$

- stabilize product dark states: exactly local
- realize competition: do not commute with H (phase fluctuations)

Continuum (1+1) dimensional Model

- model: phenomenologically motivated continuum limit for lattice fermions (weak measurement regime)

- Hamiltonian: massless Dirac fermions $\hat{\Psi}_x = (\hat{\psi}_{R,x}, \hat{\psi}_{L,x})^T$



$$\hat{H} = iv \int_x \hat{\Psi}_x^\dagger \sigma_z \partial_x \hat{\Psi}_x$$

fermionic variant

bosonization
dictionary

$$\hat{H} = \frac{v}{2\pi} \int_x [(\partial_x \hat{\theta}_x)^2 + (\partial_x \hat{\phi}_x)^2]$$

phase density
bosonized variant

- measurement operators: local particle density $\rightarrow$ current and vertex operators

rate γ_1 : $\hat{O}_{1,x} = \Psi_x^\dagger \Psi_x = \hat{J}_x^{(0)}$

$\longrightarrow$

$\hat{O}_{1,x} = -\frac{1}{\pi} \partial_x \hat{\phi}_x$ linear gapless

rate γ_2 : $\hat{O}_{2,x} = \Psi_x^\dagger \sigma_x \Psi_x$

$\longrightarrow$

$\hat{O}_{2,x} = m \cos(2\hat{\phi}_x)$ nonlinear

- $\rightarrow$ intuition: bosonic theory is long-wavelength hydrodynamics of conserved fermions
- $\rightarrow$ problem setting: **non-Hermitian Gaussian theory + cos-nonlinearity**

Towards the Relevant Degrees of Freedom: Replica Approach (n=2)

- Access state-dependent observables, e.g. covariance matrix

$$C_{xy} = \overline{\langle \hat{n}_x \hat{n}_y \rangle} - \overline{\langle \hat{n}_x \rangle} \overline{\langle \hat{n}_y \rangle} = \overline{\text{tr}[\hat{n}_x \hat{n}_y \hat{\rho}] \cdot \text{tr}[\hat{\rho}]} - \overline{\text{tr}[\hat{n}_x \hat{\rho}] \cdot \text{tr}[\hat{n}_y \hat{\rho}]}$$

- **Replica approach** via tensor products (analogous: introducing spatial index in second quantization)


- replica conditioned density matrix

$$\hat{\rho}^{(1)} \equiv \hat{\rho} \otimes \mathbf{1}, \quad \hat{\rho}^{(2)} \equiv \mathbf{1} \otimes \hat{\rho}$$

$$\hat{\rho}_2 = \hat{\rho}^{(1)} \hat{\rho}^{(2)} = \hat{\rho} \otimes \hat{\rho}$$
- replica operators, e.g.

$$\hat{n}_x^{(1)} \equiv \hat{n}_x \otimes \mathbf{1}, \quad \hat{n}_x^{(2)} \equiv \mathbf{1} \otimes \hat{n}_x$$

$$\hat{\rho} = |\psi\rangle\langle\psi|$$

$$|\Psi\rangle = |\psi^{(1)}\rangle|\psi^{(2)}\rangle =$$


- Covariance matrix:

$$C_{xy} = \frac{1}{2} \overline{\text{tr}[(\hat{n}_x^{(1)} - \hat{n}_x^{(2)})(\hat{n}_y^{(1)} - \hat{n}_y^{(2)})\hat{\rho}_2]} = \frac{1}{2} \text{tr}[(\hat{n}_x^{(1)} - \hat{n}_x^{(2)})(\hat{n}_y^{(1)} - \hat{n}_y^{(2)})\overline{\hat{\rho}_2}]$$

- More generally: all quadratic-in-state observables encoded in

$$\overline{\hat{\rho}_2} = \overline{|\Psi_t\rangle\langle\Psi_t|} \quad \Rightarrow \text{linear statistical average of replica density matrix}$$



Towards the Relevant Degrees of Freedom: Replica Approach (n=2)

- Goal: derive evolution equation for 2-replica density matrix
 - ➔ can be decoupled for Gaussian theory
 - ➔ interacting theory: useful truncation possible
 - ➔ fits problem well: impact of weak non-linear term on top of Gaussian theory. E.g. density covariance:

$$C_{xy} = \frac{1}{2} \text{tr}[(\partial_x \hat{\phi}_x^{(1)} - \partial_x \hat{\phi}_x^{(2)})(\partial_y \hat{\phi}_y^{(1)} - \partial_y \hat{\phi}_y^{(2)}) \overline{\hat{\rho}_2}] \quad \text{Gaussian correlator}$$

- replicated stochastic quantum master equation (n = 2)

$$\partial_t \hat{\rho}_2 =$$


+


$$\sum_{m=1}^2 \left[-i[\hat{H}^{(m)}, \hat{\rho}_2] - \frac{\gamma}{2} \sum_l [\hat{M}_l^{(m)}, [\hat{M}_l^{(m)}, \hat{\rho}_2]] \right] + \gamma \sum_l \{ \hat{M}_l^{(1)}, \{ \hat{M}_l^{(2)}, \hat{\rho}_2 \} \}$$

individual heating Lindbladians

replica coupling

$$+ dW \sum_{m=1}^2 \sum_l \{ \hat{M}_l^{(m)}, \hat{\rho}_2 \} \quad \text{stochastic forcing}$$




- decoupling for Gaussian theories

Boson Replica Quantum Master Equation: Free Theory

- Boson measurement $H^{(\alpha)} = \frac{\nu}{2\pi} \int_x (\partial_x \hat{\phi}^{(\alpha)})^2 + (\partial_x \hat{\theta}^{(\alpha)})^2$

$$\hat{O}_{1,x}^{(\alpha)} = -\frac{1}{\pi} \partial_x \hat{\phi}_x^{(\alpha)} \quad \hat{O}_{2,x}^{(\alpha)} = 0 \quad \text{linear case first (Gaussian theory)}$$

- New degrees of freedom



$$\begin{aligned} & : \hat{\phi}^{(a)} = \hat{\phi}^{(1)} + \hat{\phi}^{(2)} \quad \text{average coordinate} \\ & : \hat{\phi}^{(r)} = \hat{\phi}^{(1)} - \hat{\phi}^{(2)} \quad \text{replica fluctuations} \end{aligned}$$

→ Master equation becomes **separable**

- Average coordinate: **heating** to infinite temperature ($\longleftrightarrow$ unbounded growth of mode occupation)

$$\partial_t \rho^{(a)} = i[\rho^{(a)}, H^{(a)}] + \frac{2\gamma}{\pi} \sum_l \left(\partial_x \hat{\phi}^{(a)} - \overline{\langle \partial_x \hat{\phi}^{(a)} \rangle} \right) \rho^{(a)} \left(\partial_x \hat{\phi}^{(a)} - \overline{\langle \partial_x \hat{\phi}^{(a)} \rangle} \right) \leftarrow \text{only jump term!}$$

- Relative coordinate: **cooling/damping** into dark state

$$\partial_t \rho^{(r)} = i[\rho^{(r)}, H^{(r)}] - \frac{\gamma}{\pi} \sum_l \left\{ (\partial_x \hat{\phi}^{(r)})^2, \rho^{(r)} \right\} \leftarrow \text{no jump term!}$$

- further separable: $\partial_t |\psi_t^{(r)}\rangle = -i H_{\text{eff}} |\psi_t^{(r)}\rangle$

$$H_{\text{eff}} = \frac{\nu}{2\pi} \int_x (\partial_x \hat{\theta})^2 + (1 - i\eta^2) (\partial_x \hat{\phi})^2$$

$$\eta^2 = \frac{2\gamma}{\nu}$$

gapless non-Hermitean Hamiltonian

- observable of interest exclusively depends on relative coordinate


$$C_{xy} = \frac{1}{2} \text{tr} [\partial_x \hat{\phi}_x^{(r)} \partial_y \hat{\phi}_y^{(r)} \overline{\hat{\rho}_2}]$$

Boson Replica Quantum Master Equation: Including Interactions

- Boson measurement $H^{(\alpha)} = \frac{\nu}{2\pi} \int_x (\partial_x \hat{\phi}^{(\alpha)})^2 + (\partial_x \hat{\theta}^{(\alpha)})^2$

$$\hat{O}_{1,x}^{(\alpha)} = -\frac{1}{\pi} \partial_x \hat{\phi}_x^{(\alpha)} \quad \hat{O}_{2,x}^{(\alpha)} = m \cos(2\hat{\phi}_x) \quad \text{general case}$$

- $\hat{O}_{2,x}^{(\alpha)}$ couples relative and absolute degrees $\sim m \cos(\sqrt{2}(\hat{\phi}_x^{(a)} \pm \hat{\phi}_x^{(r)}))$

→ how do  degrees enter? → $\langle \hat{\phi}_x^{(a)} \hat{\phi}_x^{(a)} \rangle = \infty$ nonlinearity irrelevant for $\hat{\phi}_x^{(a)}$



→ integrate out $\hat{\phi}_x^{(a)}$ in Gaussian approx. for $\hat{\rho}^{(a)}$

- Non-hermitian Schrödinger equation for relative coordinate

$$\partial_t |\psi_t^{(r)}\rangle = -i H_{\text{eff}} |\psi_t^{(r)}\rangle \quad \rightarrow \text{cooling into dark state}$$

$$H_{\text{eff}} = \frac{\nu}{2\pi} \int_x (\partial_x \hat{\theta})^2 + (1 - i\eta^2) (\partial_x \hat{\phi})^2 - i \frac{\gamma m}{\pi} \int_x [1 - \cos(\sqrt{8} \hat{\phi}_x)]$$

effect of non-linearity

→ non-Hermitian Sine-Gordon: pinning via cos term, depinning via theta term

→ extract physics in path integral approach

Effective Non-Hermitian Hamiltonian and Path Integral

➔ Sine-Gordon action with complex coefficients

Fendley, Saleur, Zamolodchikov,
International Journal of Modern Physics (1993)

$$S = \int_{t,x} \left\{ \frac{K}{16\pi} \left[\frac{1}{\eta} (\partial_t \phi)^2 - \eta (\partial_x \phi)^2 \right] - i\lambda \cos(\phi) \right\}$$



• ‘Wick rotation’ brings free part to standard Euclidean (2+0) dimensional form $(x, t) \rightarrow (\eta^{\frac{1}{2}} x, i\eta^{-\frac{1}{2}} t)$

➔ RG flow: standard KT flow with complex K, λ

$$\partial_s \lambda = \left(2 - \frac{8\pi}{K} \right) \lambda,$$

$$\partial_s K = -\lambda^2$$

- UV flow modified
- ➔ shift of phase border
- IR flow reaches standard KT flow
- ➔ same long wavelength properties



interaction irrelevant:
flow to **free** gapless
theory -> Gaussian
approximation good

interaction relevant:
flow to free massive
theory

- ➔ gapless generalized CFT phase with algebraic correlations and varying exponent
- ➔ phase transition in the BKT universality class in line with numerics

n-Replicas: Lindblad-Keldysh 2.0


n-replica Hamilton-Keldysh: Aleiner, Faoro, Ioffe, AoP (2016); Tsuji, Werner, Ueda, PRA (2017); Shenker, Stanford, JHEP (2015); Ansari, Nazarov, JETP (2016)

- Motivation:
 - Generalization of ‘hot’ and ‘cold’ modes?
 - Entanglement entropies?

- Lindblad-Keldysh construction for n replicas

- evolution operator $\hat{V}_{dt} = \exp \left[-(i\hat{H} + \hat{M}_t^2)dt + \xi \hat{M}_t \right]$ expansion to second order: SSE

- single replica



n-Replicas: Lindblad-Keldysh 2.0

n-replica Hamilton-Keldysh: Aleiner, Faoro, Ioffe, AoP (2016); Tsuji, Werner, Ueda, PRA (2017); Shenker, Stanford, JHEP (2015); Ansari, Nazarov, JETP (2016)

- Motivation:


- Generalization of ‘hot’ and ‘cold’ modes?
- Entanglement entropies?

- Lindblad-Keldysh construction for n replicas

- evolution operator $\hat{V}_{dt} = \exp \left[-(i\hat{H} + \hat{M}_t^2)dt + \xi \hat{M}_t \right]$ expansion to second order: SSE

- single replica

$$Z(1, \{\xi\}) = \text{tr} \rho_{t_f \rightarrow \infty} =$$



$$\overline{(\dots)} \quad \overline{Z(1, \{\xi\})} \equiv Z(1)$$

$$Z(1, \{\xi\}) = \int \mathcal{D}[\Psi] \exp [i(S_{1,H}[\Psi] + S_{1,\xi}[\Psi])]$$

$$\implies Z(1) = \int \mathcal{D}\Psi \exp [i(S_{1,H}[\Psi] + S_{1,M}[\Psi])]$$

➔ measurement expectation values cancel

➔ Lindblad-Keldysh functional integral reproduced (herm. Lindblads)

$$S_{1,H}[\Psi] = \sum_{\sigma=\pm} \sigma \int_t (\bar{\psi}_\sigma i \partial_t \psi_\sigma - H[\bar{\psi}_\sigma, \psi_\sigma])$$

$$S_{1,\xi}[\Psi] = i \sum_{\sigma=\pm} \int_t [M_\sigma^2 - \xi M_\sigma]$$

$$S_{1,M}[\Psi] = i \int_t [M_+^2 + M_-^2 - \frac{1}{2}(M_+ + M_-)^2]$$

$$= -i \int_t [O_+ O_- - \frac{1}{2} O_-^2 + \frac{1}{2} O_+^2]$$

$$M = O - \bar{O}$$

n-Replicas: Lindblad-Keldysh 2.0

n-replica Hamilton-Keldysh: Aleiner, Faoro, Ioffe, AoP (2016); Tsuji, Werner, Ueda, PRA (2017); Shenker, Stanford, JHEP (2015); Ansari, Nazarov, JETP (2016)


- Motivation:

- Generalization of ‘hot’ and ‘cold’ modes?
- Entanglement entropies?

- Lindblad-Keldysh construction for n replicas

- evolution operator $\hat{V}_{dt} = \exp \left[-(i\hat{H} + \hat{M}_t^2)dt + \xi \hat{M}_t \right]$ expansion to second order: SSE

- n replicas



$$Z(n, \{\xi\}) = \int \mathcal{D}[\Psi] \exp [i(S_{n,H}[\Psi] + S_{n,\xi}[\Psi])]$$

$$\Rightarrow Z(n) = \int \mathcal{D}\Psi \exp [i(S_{n,H}[\Psi] + S_{n,M}[\Psi])]$$

➡ collective coupling to noise

➡ structural simplification for linear measurement dynamics


$$S_{n,H}[\Psi] = \sum_{\sigma=\pm} \sum_{l=1}^n \sigma \int_t \left(\bar{\psi}_{\sigma}^{(l)} i \partial_t \psi_{\sigma}^{(l)} - H[\bar{\psi}_{\sigma}^{(l)}, \psi_{\sigma}^{(l)}] \right)$$

$$S_{n,\xi}[\Psi] = i \sum_{\sigma=\pm} \sum_{l=1}^n \int_t \left[(M_{\sigma}^{(l)})^2 - \xi M_{\sigma}^{(l)} \right]$$

$$S_{n,M}[\Psi] = i \int_t \sum_{l=1}^n \left[\left([M_{+}^{(l)}]^2 + [M_{-}^{(l)}]^2 \right) \right. \\ \left. - \frac{1}{2} \left(\sum_{l=1}^n M_{+}^{(l)} + M_{-}^{(l)} \right)^2 \right]$$

collective coupling
to noise!

n-Replicas: Decoupling of Gaussian theories



- practical importance: reduction to linear / Gaussian bosonic theory in limiting cases

- bosonized action in the presence of noise:

$$S[\phi] = S_{n,H}[\phi] + S_{n,\xi}[\phi]$$

bosonic field

$$S_{n,H}[\phi] = -\frac{1}{2\pi} \sum_{l=1}^n \sum_{\sigma=\pm} \sigma \int_{t,x} \phi_{\sigma}^{(l)} (\partial_t^2 - \partial_x^2) \phi_{\sigma}^{(l)} \quad S_{n,\xi}[\phi] = i \sum_{\sigma=\pm} \sum_{l=1}^n \int_{t,x} \left[(O_{\sigma}^{(l)} - \bar{O})^2 - \xi (O_{\sigma}^{(l)} - \bar{O}) \right]$$

$$O_{\sigma}^{(l)} = D \phi_{\sigma}^{(l)}, \quad D = \pi m, \partial_x$$

quadratic

- **decoupling** of center-of-mass and relative modes

- Fourier expansion in replica space

$$\phi_{\sigma,t,x}^{(l)} = \frac{1}{\sqrt{n}} \sum_{k=0}^{n-1} e^{-i \frac{2\pi k l}{n}} \phi_{\sigma,t,x}^{(k)}$$

- equation of motion $\frac{\delta S}{\delta \phi} = 0$

$$\partial_t^2 \phi_{+}^{(k)} = (\partial_x^2 - \frac{\gamma i}{\pi v} D^2) \phi_{+}^{(k)} + \sqrt{n} \xi \delta_{k,0}$$


➔ no reference to measurement expectation value

➔ exact decoupling into:


➔ 1 collective 'hot' mode, heating to infinite temperature

➔ (n-1) 'cold' modes, do not 'see' the noise, cool to ground state of non-hermitean Hamiltonian

replica space picture




reciprocal replica space picture



Entanglement Entropies from Replica Approach

- Keldysh path integral representation of n-th moment $Z_A(n, \{\xi\}) = \text{tr} \rho_A^n$



- formulate boundary condition via operator insertion (in the presence of noise):

$$Z_A(n, \{\xi\}) = \int \mathcal{D}[\Psi] T e^{iS_n[\Psi]} \quad T\Psi_{+,x,t_f}^{(l)} = \begin{cases} -\Psi_{-,x,t_f}^{(l)} & \forall x \notin A \\ -\Psi_{-,x,t_f}^{(l+1)} & \forall x \in A \end{cases}$$

translation by one in replica space

- e.g. free massless Dirac fermions (after bosonization into equivalent Luttinger Hamiltonian): factorization


$$Z_A(n, \{\xi\}) = \left\langle \prod_{k=1}^n \exp \left(-\sqrt{2}i \frac{k}{n} \int dx [\delta(x - x_0) - \delta(x - (x_0 + L))] \phi_{c,x,t}^{(k)} \right) \right\rangle$$

ground states: Casini, Fosco, Huerta, J. Stat. Mech. (2005)

- ➡ boundary conditions appear as opposite charges
- ➡ $k = 0$ mode does not contribute!
- ➡ $k \neq 0$ independent of noise ξ !

Entanglement Entropies from Replica Approach

- Keldysh path integral representation of n-th moment $Z_A(n, \{\xi\}) = \text{tr} \rho_A^n$



- n-th Rényi entropy

$$S_n = \frac{1}{1-n} \overline{\log Z_A(n, \{\xi\})}$$


- von Neumann entropy

$$S = \lim_{n \rightarrow 1} S_n = \frac{2}{3} \langle \phi_{c,x_0}^{(k>0)} \phi_{c,x_0+L}^{(k>0)} \rangle$$



➡ correlator in the Gaussian dark state wave function

Construction summary and entanglement entropy

- Rényi entropy $S_n(L) = \frac{1}{1-n} \overline{\log Z_A(n, \{dW\})}$, $Z_A(n, \{dW\}) \equiv \text{tr}[(\hat{\rho}_A^{(c)})^n]$
- von Neumann entropy: $n \rightarrow 1$
- in practice need theory for n Keldysh replicas



- **decoupling** of center-of-mass and relative modes (**exact** for Gaussian states -> good away from transition)



- observation 1:

- 1 mode heats up (noisy)
- n-1 modes cool down (noiseless)


- observation 2:

- noisy contribution A independent
- all A dependence in noiseless modes!

- ➔ entropy determined by noiseless modes
- ➔ rationalizes analogy to ground state phase transition

Entanglement Transition from Replica Approach

- focus on von Neumann entropy S in Gaussian limiting regimes




$$S = \frac{1}{3}c(\gamma) \log(L)$$

$$S = \frac{1}{3}c(\gamma) \log(m^{-1}) \sim L^0$$


- sub-volume log-law
- $c(\gamma \rightarrow 0) \rightarrow 1$
 - ➔ ground state entropy of massless Dirac
- in Gaussian state:
 - $c(\gamma \rightarrow \infty) \sim \gamma^{-1/2}$
 - ➔ compatible with numerics in critical phase
- with RG improvement, qualitatively similar to numerics

- saturation to area law



Entanglement Transition from Replica Approach

- focus on von Neumann entropy S in Gaussian limiting regimes




$$S = \frac{1}{3}c(\gamma) \log(L)$$

$$S = \frac{1}{3}c(\gamma) \log(m^{-1}) \sim L^0$$


- sub-volume log-law
- $c(\gamma \rightarrow 0) \rightarrow 1$
 - ➔ ground state entropy of massless Dirac
- in Gaussian state:
 - $c(\gamma \rightarrow \infty) \sim \gamma^{-1/2}$
 - ➔ compatible with numerics in critical phase
- non-commuting limit: $\gamma = 0$
finite temperature initial state
 - $S \sim L$
 - ➔ volume law $\longleftrightarrow$ finite temperature massless Dirac

- saturation to area law




- ➔ underpins entanglement transition at finite critical g
- ➔ picture qualitatively in line with numerics

Revealing measurement-induced phase transitions by pre-selection



Thomas Müller

M. Buchhold, T. Müller, SD,
arxiv:2208.10506 (2022)




Michael Buchhold

Observability of the transition?

- Standard quantum mechanical observables: need to reproduce **identical** state
- Randomness of measurements outcomes masks transition in qm observables

- Post-selection problem: [way out for Cliffords: Gullans, Huse, PRL \(2020\);](#)
[exp: C. Noel et al. Nat Phys. \(2021\)](#)



- Select one string of measurement outcomes (trajectory)
- build an ensemble of identical states with same trajectory

➔ number of outcomes (single site)

$2^{\#}$ of measurements


➔ post-selection **exponentially costly**

- Pre-selection solution: [M. Buchhold, T. Müller, SD, arxiv:2208.10506 \(2022\)](#)

- break randomness in measurement outcomes
- steer system into unique representative state in Hilbert space (**dark state**)

➔ can be done **gently**: no modification of entanglement structure & universal properties (as post-selection)

➔ **no exponential overhead**: observable by standard means in NISQ platforms



Pre-selection concept


- pull MIPT to observable level $\longleftrightarrow$ study measurement averaged density matrix

$$\hat{\rho}_{t+\delta t} = \hat{\mathcal{X}}[\hat{\rho}_t]\delta t \quad \text{e.g. temporal continuum limit} \quad \hat{\mathcal{X}}[\hat{\rho}] = -i[\hat{H}, \hat{\rho}] + \sum_l \hat{L}_l \hat{\rho} \hat{L}_l^\dagger - \frac{1}{2} \{ \hat{L}_l^\dagger \hat{L}_l, \hat{\rho} \}$$

- unmodified dynamics w/o preselection:

$$\hat{\mathcal{X}}^{(0)} \quad \text{with} \quad \hat{H}^{(0)}, \quad \hat{L}_l^{(0)} = (\hat{L}_l^{(0)})^\dagger \quad \text{hermitian}$$

$$\implies \hat{\mathcal{X}}^{(0)}[\hat{\mathbf{1}}] = 0 \quad \text{fully mixed stationary state}$$



- modified dynamics with pre-selection:

$$\hat{H}^{(0)}, \hat{L}^{(0)} \rightarrow \hat{H}, \hat{L} \quad \text{such, that}$$

1. There exists a **pure dark state** representing one possible measurement outcome,

$$\hat{\rho}_D = |D\rangle\langle D| \quad \text{s.t.} \quad \hat{\mathcal{X}}[\hat{\rho}_D] = 0$$


2. The dark state is **unique**

3. Modifications preserve **symmetries** and are **irrelevant in the RG sense**

- $\Rightarrow$ pre-select representative state (e.g. with order parameter)

- $\Rightarrow$ directionality in Hilbert space

- $\Rightarrow$ phases and phase transitions unmodified




Pre-selection: connection to absorbing state transitions

- dynamics $\partial_t \hat{\rho} = \mathcal{X}(\hat{\rho}) = \mathcal{X}_1(\hat{\rho}) + \gamma \mathcal{X}_2(\hat{\rho})$
unitary non-unitary


$\mathcal{X}_1(\hat{\rho}) :$

$\mathcal{X}_2(\hat{\rho}) :$

prefers
entangled



prefers
product



trajectory entanglement entropy:

$$\gamma \geq \gamma_c : S(\hat{\rho}_A) \sim |A|^0$$

$$\gamma < \gamma_c : S(\hat{\rho}_A) \sim |A|, \log(|A|)$$

but: extensive configuration entropy

- unmodified measurement dynamics:

Pre-selection: connection to absorbing state transitions


- dynamics $\partial_t \hat{\rho} = \mathcal{X}(\hat{\rho}) = \mathcal{X}_1(\hat{\rho}) + \gamma \mathcal{X}_2(\hat{\rho})$

unitary
non-unitary


$\mathcal{X}_1(\hat{\rho}) :$

$\mathcal{X}_2(\hat{\rho}) :$

prefers
entangled



prefers
product




trajectory entanglement entropy:

$$\gamma \geq \gamma_c : S(\hat{\rho}_A) \sim |A|^0$$

$$\gamma < \gamma_c : S(\hat{\rho}_A) \sim |A|, \log(|A|)$$


but: extensive configuration entropy

dynamically
unstable



dynamically
stable

$\hat{\rho}_D$



phase transition:

pre-selection fails

$$\gamma < \gamma_c : \hat{\rho} \not\rightarrow \hat{\rho}_D$$

pre-selection succeeds

$$\gamma \geq \gamma_c : \hat{\rho} \rightarrow \hat{\rho}_D$$

configuration entropy removed!

Pre-selection: connection to (quantum) absorbing state transitions


- dynamics $\partial_t \hat{\rho} = \mathcal{X}(\hat{\rho}) = \mathcal{X}_1(\hat{\rho}) + \gamma \mathcal{X}_2(\hat{\rho})$

unitary
non-unitary

$\mathcal{X}_1(\hat{\rho}) :$

$\mathcal{X}_2(\hat{\rho}) :$


prefers
entangled



product
states

entangled
states

prefers
product




trajectory entanglement entropy:

$$\gamma \geq \gamma_c : S(\hat{\rho}_A) \sim |A|^0$$


$$\gamma < \gamma_c : S(\hat{\rho}_A) \sim |A|, \log(|A|)$$

but: extensive configuration entropy

dynamically
unstable




dynamically
stable



$\hat{\rho}_D$

(up to time scale exp in system size)



wetting
transition

wetted/active

critical

absorbing/dry

$$\gamma \geq \gamma_c : \hat{\rho} \rightarrow \hat{\rho}_D$$

$$\gamma < \gamma_c : \hat{\rho} \not\rightarrow \hat{\rho}_D$$


→ transition in the dynamics

→ structure of absorbing state
transition in Hilbert space!

H. Hinrichsen, Adv. Phys. (2000);
G. Odor, RMP (2004)

Example: 'Classical' preselection dynamics in monitored fermions

- choice of representative dark state: $\hat{\rho}_D = |101010\dots\rangle\langle 101010\dots|$




- modify generator of dynamics: condition hopping on state (non-linear in state):

$$\hat{H}^{(0)} \rightarrow \hat{H} = \hat{H}[\hat{\rho}] \quad \text{with} \quad \hat{H}[\hat{\rho}_D] = 0$$

Wiseman, Milburn, PRL (1993)


$$J \rightarrow J_l[\hat{\rho}] = 2 - (\lfloor \langle \hat{n}_{l-1} \rangle \rfloor - \lfloor \langle \hat{n}_l \rangle \rfloor)^2 - (\lfloor \langle \hat{n}_{l+1} \rangle \rfloor - \lfloor \langle \hat{n}_{l+2} \rangle \rfloor)^2$$

- distinguishes phases via Néel order



Example: 'Classical' preselection dynamics in monitored fermions

- choice of representative dark state: $\hat{\rho}_D = |101010\dots\rangle\langle 101010\dots|$




- modify generator of dynamics: condition hopping on state (non-linear in state):

$$\hat{H}^{(0)} \rightarrow \hat{H} = \hat{H}[\hat{\rho}] \quad \text{with} \quad \hat{H}[\hat{\rho}_D] = 0$$


Wiseman, Milburn, PRL (1993)

$$J \rightarrow J_l[\hat{\rho}] = 2 - (\lfloor \langle \hat{n}_{l-1} \rangle \rfloor - \lfloor \langle \hat{n}_l \rangle \rfloor)^2 - (\lfloor \langle \hat{n}_{l+1} \rangle \rfloor - \lfloor \langle \hat{n}_{l+2} \rangle \rfloor)^2$$

- distinguishes phases via Néel order



- dark state phase transition: BKT



- entanglement entropies coincide: phases / phase transition unmodified
- but there exist standard qm observables witnessing it

Example: 'quantum' preselection

- modify operators: linear in state

$$\partial_t \rho = i[\rho, H] - \gamma \sum_l \{L_l^\dagger L_l, \rho\} - 2L_l \rho L_l^\dagger$$

modified measurement operators

$$[\hat{H}, \hat{\rho}_D] = 0$$




$$\hat{L}_l \hat{\rho}_D = 0$$

- measurement preselection:

$$\hat{L}_l = \hat{U}_l \hat{n}_l$$


$$[\hat{U}_l, \hat{n}_l] \neq 0$$

- non-Gaussian evolution



DMRG simulations, M. Buchhold, T. Müller, SD, in progress

- detect transition



- field theory

$$\hat{L}_l = \hat{n}_l \rightarrow \exp[i\pi(c_l^\dagger c_m + c_m^\dagger c_l)] \hat{n}_l$$

$$\partial_x \hat{\phi}_x + \cos(\hat{\phi}_s) \rightarrow \partial_x \hat{\phi}_x + \cos(\hat{\phi}_s) + i\partial_x \hat{\theta}_x$$

- Heisenberg-Langevin theory (or equivalent Keldysh)

$$(\partial_t^2 - \partial_x^2) \hat{\phi}_x + \gamma \cos \hat{\phi}_x + \gamma \sin \hat{\phi}_x (\partial_t \hat{\phi}_x + \sin \hat{\phi}_x) = \hat{\xi}_x$$


- ➔ irrelevant in weak measurement regime: infinite T state
- ➔ becomes relevant at transition: pinning

Outlook: Quantum absorbing transitions beyond directed percolation?

- Grassberger-Janssen conjecture: No!

Janssen, Z. Phys. B (1981); Grassberger, Z. Phys. B (1982)

- Qualitative overview:



- possible pure state scenarios:


Sieberer, Buchhold, SD, ROPP (2016)

- general quantum dynamics w/ dark/absorbing state (or equivalent Keldysh)

$$\partial_t \hat{\rho} = -i[\hat{H}, \hat{\rho}] + \gamma \sum_i \hat{L}_i \hat{\rho} \hat{L}_i^\dagger - \frac{1}{2} \{ \hat{L}_i^\dagger \hat{L}_i, \hat{\rho} \} \quad \hat{H} = \sum_i \hat{h}_i \rightarrow \hat{H} = \sum_i \hat{h}_i \hat{L}_{i-1}^\dagger \hat{L}_{i-1}$$

repulsive fixed point

attractive fixed point



dark/absorbing state

$$\hat{L}_i |D\rangle = 0 \quad \forall i$$

Outlook: Quantum absorbing transitions beyond directed percolation?

- Grassberger-Janssen conjecture: No! Janssen, Z. Phys. B (1981); Grassberger, Z. Phys. B (1982)
- Qualitative overview:

classical

mixed mixed

quantum

pure pure

quantum absorbing

mixed pure
- possible pure state scenarios: Sieberer, Buchhold, Diehl ROPP (2016)
 - general quantum dynamics w/ dark/absorbing state (or equivalent Lindblad-Keldysh functional integral)

$$\partial_t \hat{\rho} = -i[\hat{H}, \hat{\rho}] + \gamma \sum_i \hat{L}_i \hat{\rho} \hat{L}_i^\dagger - \frac{1}{2} \{ \hat{L}_i^\dagger \hat{L}_i, \hat{\rho} \} \qquad \hat{H} = \sum_i \hat{h}_i \rightarrow \hat{H} = \sum_i \hat{h}_i \hat{L}_{i-1}^\dagger \hat{L}_{i-1}$$


$$\hat{L}_i |D\rangle = 0 \quad \forall i \quad \text{dark/absorbing state}$$

		State	
		product (w/o entanglement)	entangled ('non-trivial vacua')
Dynamics	w/o conservation law	directed percolation	$ D\rangle =$ (BCS superfluid)
	w/ conservation law	1+1: present problem, BKT d+1, d > 1: tbd	$ D\rangle =$ (topological insulator)

Summary


General non-unitary quantum dynamics hosts new types of phases and phase transitions

- here: critical to area law entanglement phase transition
- BKT transition revealed by Keldysh replica field theory



Observability via pre-selection

- like post-selection, but no exponential overhead
- various strategies for NISQ platforms
- link to quantum absorbing state transitions



Outlook

- general principles for observability?
- higher dimensions: novel quantum absorbing state transitions beyond directed percolation?
- relation of measurement vs. disorder problems? role of conservation laws? symmetry classification? [Poboiko, Pöpperl, Gornyi, Mirlin, arXiv:2304.03138 \(2023\)](#)
- emergent CFT behavior?

