University of Cologne

http://www.thp.uni-koeln.de/gravitation/courses/rci2223.html

Institute for Theoretical Physics

Prof. Dr. Claus Kiefer

Tatevik Vardanyan, Mauricio Ortiz Torres and Blanca Hergueta

ver. 1.01

Third exercise sheet on Relativity and Cosmology I

Winter term 2022/23

Release: Thu, Nov. 3rd

Submit: Thu, Nov. 10th

Discuss: Thu, Nov. 17th

Exercise 6 (10 points): Covariant Maxwell equations

Recall from classical electromagnetism the Maxwell equations (in Gaussian units with c = 1):

$$\vec{\nabla} \cdot \vec{E} = 4\pi\rho, \quad \vec{\nabla} \times \vec{B} - \frac{\partial \vec{E}}{\partial t} = 4\pi \vec{J}; \qquad \vec{\nabla} \cdot \vec{B} = 0, \quad \vec{\nabla} \times \vec{E} + \frac{\partial \vec{B}}{\partial t} = 0.$$
 (1)

In terms of the scalar (Φ) and vector (\vec{A}) potentials, the electric and magnetic fields are $\vec{E} = -\vec{\nabla}\Phi - \hat{c}_t\vec{A}$ and $\vec{B} = \vec{\nabla} \times \vec{A}$. In components, eq. (1) reads

$$\sum_{i=1}^{3} \partial_{i} E^{i} \equiv \partial_{i} E^{i} = 4\pi\rho, \qquad \qquad \sum_{j=1}^{3} \sum_{k=1}^{3} \epsilon^{ijk} \partial_{j} B_{k} - \partial_{t} E^{i} \equiv \epsilon^{ijk} \partial_{j} B_{k} - \partial_{t} E^{i} = 4\pi J^{i}; \qquad (2a)$$

$$\partial_i B^i = 0$$
, $\epsilon^{ijk} \partial_j E_k + \partial_t B^i = 0$, (2b)

where e^{ijk} is the Levi-Civita pseudo-tensor, $i, j, \ldots = 1, 2, 3$, and Einstein notation is assumed, as has been explained in eq. (2a). Let $A^{\mu} := (\Phi, \vec{A})$ be the four-potential, $j^{\mu} := (\rho, \vec{J})$ the four-current, $\mu, \nu, \ldots = 0, 1, 2, 3$, and define the field-strength tensor

$$F_{\mu\nu} := \partial_{\mu} A_{\nu} - \partial_{\nu} A_{\mu} \,. \tag{3}$$

6.1 Show that the expressions

$$\sum_{\nu=0}^{4} \partial_{\nu} F^{\mu\nu} \equiv \partial_{\nu} F^{\mu\nu} = 4\pi j^{\mu} \,, \tag{4a}$$

$$\partial_{\rho}F_{\mu\nu} + \partial_{\mu}F_{\nu\rho} + \partial_{\nu}F_{\rho\mu} = 0 \tag{4b}$$

correspond to eqs. (2a) and (2b) respectively.

Hint: $\epsilon_{ijk}\epsilon^{ilm} = \delta_i^l \delta_k^m - \delta_i^m \delta_k^l$.

6.2 Show that eq. (4a) leads to the continuity equation $0 = \partial_t \rho + \vec{\nabla} \cdot \vec{J} \equiv \partial_\mu j^\mu$. How does the continuity equation look like in a Lorentz-boosted reference frame?

Exercise 7 (6 points): Covariant Lorentz force

Let $p^{\mu} := mu^{\mu}$ be the kinematic four-momentum, u^{μ} the four-velocity, τ the proper time; $\vec{P} := m\vec{v}$, $\vec{v} := d\vec{x}/dt$, and t the coordinate time.

7.1 Show that the spatial components of the covariant Lorentz four-force

$$\frac{\mathrm{d}p_{\mu}}{\mathrm{d}\tau} = f_{\mu} = qF_{\mu\nu}u^{\nu} \tag{5}$$

give in the non-relativistic limit the Lorentz force

$$\frac{\mathrm{d}\vec{P}}{\mathrm{d}t} = \vec{F} = q(\vec{E} + \vec{v} \times \vec{B}).$$

7.2 What is the physical meaning of the time component f^0 of the covariant four-force in eq. (5)?

Exercise 8 (6 points): *Kottler–Møller coordinates*

Let g be a constant acceleration, (t, x, y, z) Cartesian coordinates in 4-dimensional Minkowski space. Consider the transformation (c = 1)

$$t = \left(\frac{1}{g} + x'\right) \sinh gt',$$

$$x = \left(\frac{1}{g} + x'\right) \cosh gt' - \frac{1}{g},$$

$$y = y', \qquad z = z'.$$

- **9.1** Show that the transformation gives accelerated frames of reference with constant acceleration.
- **9.2** Calculate the components of the metric with respect to the coordinates (t', x', y', z').